

This article was downloaded by: [York University Libraries]

On: 01 February 2015, At: 22:42

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH,  
UK



## Annals and Magazine of Natural History: Series 7

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tnah13>

### VIII.—Revised Nomenclature of the Species described in Bate and Westwood's 'British Sessile-eyed Crustacea'

Canon A. M. Norman M.A., D.C.L., LL.D., F.R.S.

Published online: 01 Dec 2009.

To cite this article: Canon A. M. Norman M.A., D.C.L., LL.D., F.R.S. (1905)

VIII.—Revised Nomenclature of the Species described in Bate and Westwood's 'British Sessile-eyed Crustacea', *Annals and Magazine of Natural History: Series 7*, 16:91, 78-95, DOI: [10.1080/03745480509443653](https://doi.org/10.1080/03745480509443653)

To link to this article: <http://dx.doi.org/10.1080/03745480509443653>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

VIII.—*Revised Nomenclature of the Species described in Bate and Westwood's 'British Sessile-eyed Crustacea.'* By Canon A. M. NORMAN, M.A., D.C.L., LL.D., F.R.S.

THE work of Bate and Westwood from the first presented great difficulties to the student of the Amphipoda and Isopoda in consequence of the unsatisfactory figures and inadequate descriptions of many of the species. Moreover, with increasing knowledge of those Crustacea extensive changes in nomenclature have become necessary. Hence there lies a heavy stumbling-block in the way of any carcinologist who commences the study of the species represented in our fauna, and only an adequate knowledge of all that during recent years has been written on the subject can at the present time enable the species described in this work to be correctly named.

The object of the present paper is to remove, at any rate partially, this difficulty and bring the nomenclature of Bate and Westwood into juxtaposition with the names which are now employed for the several species. I have, of course, availed myself of all that has been written on the subject by carcinologists, such as the Rev. T. R. R. Stebbing, Mr. A. O. Walker, and others in our own country, and by Professor G. O. Sars and M. Chevreux and others on the Continent, and I trust that I shall moreover be able to throw fresh light on the relationship of certain species which have hitherto not received full elucidation.

There have been very large additions to our knowledge of the species of these two orders which inhabit our seas during recent years, but with such additions these notes have nothing to do.

One new genus is here introduced—*Coremapus* (*κόρημα*, a brush, and *πόντος*, a foot). Allied to *Microdeutopus*; first gnathopods the largest, complexly subchelate in male and simply subchelate in female as in that genus, but the second gnathopods in both sexes narrow, scarcely subchelate, terminating in a minute nail, the limb forming a brush, the meros, carpus, and manus being densely setose, especially the meros, which is much produced over the carpus. Type, *Coremapus (Microdeutopus) versiculatus*, Bate.

In the earlier part of this paper the species are taken in the order in which they are arranged in our authors' work; the nomenclature, where necessary, is corrected and a number prefixed.

In the latter part the species with their corrected names

are arranged as in the existing leading works upon the subject, those of Professor G. O. Sars, and the affixed number is that under which the species occurs in the previously given list.

I.

The Species described in Bate and Westwood, with correction of names where such correction is required.

VOL. I.

1. *Talitrus locusta* (Linné), p. 16, = *Talitrus locusta* (Pallas).
2. *Orchestia littorea* (Montagu), p. 27.
3. — *mediterranea* (Costa), p. 31.
4. — *Deshayesii*, Audouin, p. 36, = *Orchestoidea Deshayesii* (Audouin) (Norman, Ann. & Mag. Nat. Hist. ser. 7, vol. v. p. 139).
5. *Allorchestes Nilssonii* (Rathke), p. 40, = *Hyale Nilsonni* (Rathke).
6. — *imbricatus*, Bate, p. 43, = *Hyale Lubbockiana* (Bate), ♂.
7. *Nicea Lubbockiana*, Bate, p. 47, = *Hyale Lubbockiana* (Bate), ♀.
8. *Montauka monoculoides* (Montagu), p. 54, = *Stenothoe monoculoides* (Montagu).
9. — *marina*, Bate, p. 58, = *Stenothoe marina* (Bate).
10. — *Alderi*, Bate, p. 61, = *Metopa Alderi* (Bate), ♀.
11. — *pollexiana*, Bate, p. 64, = *Metopa norvegica* (Lilljeborg) (Norman, Ann. & Mag. Nat. Hist. ser. 7, vol. vi. p. 41).
12. *Danaid dubia*, Bate, p. 68, = *Cressa dubia* (Bate).
13. *Lysianassa Costæ*, Bate and Westwood (nec Milne-Edwards), p. 74, = *Lysianax septentrionalis*, Della Valle.
14. — *Audouiniana*, Bate, p. 79, = *Perrierella Audouiniana* (Bate).
15. — *atlantica*, Bate and Westwood (? Milne-Edwards), p. 82, = *Ambasia Danielsseni*, Boeck (Norman, Ann. & Mag. Nat. Hist. ser. 7, vol. v. 1900, p. 144).
16. — *longicornis*, Bate and Westwood (nec Lucas), p. 85. The entire figure and the urosome probably taken from *Orchomene humilis* (Costa), ♂, the other dissections from *Lysianax septentrionalis*, Della Valle, ♂ (see Walker, Ann. & Mag. Nat. Hist., Feb. 1892; and Norman, *ibid.* ser. 7, vol. v. p. 142).
17. *Anonyx longicornis*, Bate, p. 91, = *Lepidepecreum longicorne* (Bate), ♂.
18. — *Edwardsi*, Bate (nec Kröyer), p. 94, = *Orchomene humilis* (A. Costa) (see Norman, Ann. & Mag. Nat. Hist. ser. 7, vol. v. 1900, p. 202).

19. *Anonyx obesus*, Bate, p. 98, = *Acidostoma obesum* (Bate).  
 20. — *denticulatus*, Bate, p. 101, = *Hippomedon denticulatus* (Bate).  
 21. — *Holbölli*, Bate and Westwood (nec Kröyer), p. 104, = *Haplonyx cicada* (Fabricius).  
 22. — *minutus*, Bate and Westwood (nec Kröyer), p. 108. Perhaps a young male of *Orchomene humilis* (Norman, Ann. & Mag. Nat. Hist. ser. 7, vol. v. p. 204).  
 23. — *plautus*, Bate and Westwood (nec Kröyer), p. 111, = *Menigrates obtusifrons*, Boeck.  
 24. — *longipes*, Bate, p. 113, = *Tryphosites longipes* (Bate), ♀.  
 25. — *ampulla*, Bate and Westwood (nec Kröyer), p. 116, = *Tryphosites longipes* (Bate), ♂.  
 26. *Callisoma crenata*, Bate, p. 120, = *Callisoma Hopei*, Costa.  
 27. *Ampelisca Gaimardi*, Bate and Westwood (nec Kröyer), p. 127, = *Ampelisca typica* (Bate), ♂.  
 28. — *Belliana* (Bate), p. 135, = *Ampelisca brevicornis* (A. Costa).  
 29. *Phoxus simplex*, Bate, p. 140, = *Phoxocephalus simplex* (Bate).  
 30. — *Holbölli*, Kröyer, p. 143, = *Phoxocephalus Holbölli* (Kröyer).  
 31. — *plumosus*, Bate and Westwood (nec Kröyer), p. 146, = *Harpinia neglecta*, G. O. Sars.  
 32. *Graia imbricata*, Bate, p. 152, = *Amathilla angulosa* (Rathke), young. I do not, however, consider *A. angulosa* to be a distinct species from *A. homari* (Fabricius).  
 33. *Westwoodilla cecula*, Bate, p. 155. }  
 34. — *hyalina*, Bate, p. 158. }  
 Both these *Westwoodille*, which are the same species, were so erroneously described that even Bate himself could not recognize his own genus or species, and the very next species is called *Ædicerus parvimanus*. The name to be used for the species should be therefore, I think, that by which the species was properly described and by which it could be recognized, and the species should be called *Halimedon parvimanus* (Bate and Westwood).  
 35. *Ædicerus parvimanus*, Bate and Westwood, p. 161, = *Halimedon parvimanus* (Bate), = *H. Müllerii*, Boeck.  
 36. *Monoculodes carinatus*, Bate, p. 165.  
 37. — *Stimpsoni*, Bate, p. 168, = *Monoculodes carinatus*, Bate.  
 38. *Kroyera arenaria*, Bate, p. 173, = *Pontocrates arenarius* (Bate).  
 39. — *altamarina*, Bate and Westwood, p. 177, = *Pontocrates altamarinus* (B. & W.).  
 40. *Amphilochus manudens*, Bate, p. 180.  
 41. *Darwinia compressa*, Bate, p. 184, = *Laphystius sturionis*, Kröyer.

42. *Sulcator arenarius*, Bate, p. 189, = *Haustorius arenarius* (Slabber).
43. *Urothoe Bairdii*, Bate, p. 193, = ♂ of *U. marinus*.
44. — *marinus*, Bate, p. 195, ♀.
45. — *brevicornis*, Bate, p. 198, ♀.
46. — *elegans*, Bate, p. 200, ♂.
- The species of *Urothoe* require further working out.
47. *Liljeborgia pallida*, Bate, p. 203, = *Liljeborgia pallida*, Bate.
48. — *shetlandica*, Bate and Westwood, p. 206, = *Cheirocratus Sundevallii* (Rathke), ♂.
49. *Phædra antiqua*, Bate, p. 209. Unknown; might be an *Apherusa*, if it were not that it has a secondary appendage to the antennæ.
50. — *Kinahani*, Bate, p. 211, = *Liljeborgia Kinahani* (Bate).
51. *Isca Montagui*, H. Milne-Edwards, p. 215.
52. *Iphimedia obesa*, Rathke, p. 219.
53. — *eblanæ*, Bate, p. 221.
54. *Otus carinatus*, Bate, p. 224, = *Odius carinatus* (Bate).
55. *Pereionotus testudo* (Montagu), p. 228.
56. *Acanthonotus Owenii*, Bate, p. 232, = *Epimeria cornigera* (Fabricius).
57. *Dexamine spinosa* (Montagu), p. 237.
58. — *tenuicornis*, Bate and Westwood (nec Rathke), p. 240, = *Dexamine thea*, Boeck.
59. — *vadlorensis*, Bate and Westwood, p. 242, = *Paratyulus vadlorensis* (B. & W.).
60. *Atylus Swammerdamii* (H. Milne-Edwards), p. 246, = *Paratyulus Swammerdamii* (H. Milne-Edwards).
61. — *gibbosus*, Bate, p. 248, = *Tritata gibbosa* (Bate).
62. — *bispinosus*, Bate, p. 250, = *Apherusa bispinosa* (Bate).
63. *Pherusa bicuspis*, Bate (nec Kröyer), p. 253, = *Apherusa cirrus* (Bate), ♂, = *Apherusa borealis*, Boeck.
64. — *fucicola*, Leach, p. 255, = *Gammarella brevicaudata*, H. Milne-Edwards, ♀ (Walker, Ann. & Mag. Nat. Hist. ser. 6, vol. vii. p. 418).
65. *Calliope læviuscula* (Kröyer), p. 259, = *Calliopius læviusculus* (Kröyer).
66. — *Ossiani*, Bate, p. 261, = *Parapleustes latipes* (M. Sars).
67. — *Fingalli*, Bate and Westwood, p. 263, = *Parapleustes latipes* (M. Sars), adult.
68. — *grandoculis*, Bate, p. 265, = *Calliopius læviusculus* (Kröyer), ♂.
69. *Eusirus helvetica*, Bate, p. 267, = *Eusirus longipes*, Boeck.

70. *Leucothoe articulosa* (Montagu), p. 271, = *Leucothoe spinicarpa* (Abildgaard).
71. — *furina* (Savigny), p. 274, =? *Leucothoe furina* (Savigny).
72. *Gossea microdeutopa*, Bate, p. 277. “Appears to be *Apherusa Jurinii*, M.-Edw.” (Walker, Ann. & Mag. Nat. Hist. ser. 6, vol. xv. 1895, p. 469).
73. *Aora gracilis*, Bate, p. 280.
74. *Stimpsonia chalifera*, Bate, p. 285, = *Stimpsonella chelifera* (Bate).
75. *Microdeutopus gryllotalpa*, Bate (nec Costa), p. 289, = *Microdeutopus damnoniensis*, Bate.
76. — *Websteri*, Bate, p. 291, = *Lembos Websteri*, Bate.
77. — *anomalus*, Rathke, ♀, p. 293. ♀ of *M. damnoniensis*, if not of *M. anomalus*.
78. — *versiculatus*, Bate, p. 295, = *Coremapus versiculatus* (Bate), ♀.
79. *Protomediea hirsutimana*, Bate, p. 298, = *Leptocheirus pilosus*, Zaddach.
80. — *Whitei*, Bate, p. 300, = *Cheirocratus Sunderalli* (Rathke), ♀.
81. *Bathypporeia pilosa*, Bate and Westwood (nec Lindström), p. 304, = *Bathypporeia Williamsonia* (Bate), = *B. norvegica*, Sars.
82. — *Robertsoni*, Bate, p. 307, ♂.
83. — *pelugica*, Bate, p. 309, ♂.
84. *Niphargus aquilex*, Schiödte, p. 315, = *Niphargus subterraneus* (Leach).
85. — *fontanus*, Bate, p. 319.
86. — *Kochianus*, Bate, p. 323.
87. *Crangonyx subterraneus*, Bate, p. 327.
88. *Gammarella brevicaudata* (H. Milne-Edwards), p. 330, ♂.
89. — *Normanni*, Bate and Westwood, p. 333, = *Gammarella brevicaudata* (H. Milne-Edwards), ♀.
90. *Melita palmata* (Montagu), p. 337.
91. — *obtusata* (Montagu), p. 341, ♂.
92. — *proxima*, Bate, p. 344, = *Melita obtusata* (Montagu), ♂.
93. — *gladiosa*, Bate, p. 346.
94. *Mara grossimana* (Montagu), p. 350.
95. *Eurystheus erythrocephalus* (Lilljeborg), p. 354, = *Gammaropsis maculatus* (Johnston), ♂.
96. — *bispinimanus*, Bate, p. 357, = *Gammaropsis maculatus* (Johnston), ♀.
97. *Amathilla Sabini* (Leach), p. 361, = *Amathilla homari* (Fabricius), var. *Amathilla angulosa* (Rathke).
98. *Gummarus marinus*, Leach, p. 370.
99. — *campyllops*, Leach, p. 375.

100. *Gammarus locusta* (Linné), p. 378.
101. —— *tenuimanus*, Bate, p. 394, =? *Mæra Batei*, Norman (see Walker, Ann. & Mag. Nat. Hist. ser. 6, vol. xv. 1895, p. 471).
102. —— *Edwardsii*, Bate, p. 396 (not as yet recognized by any other author).
103. —— *pulex* (De Geer), p. 388.
104. —— *dubius*, Johnston, p. 397, =?
105. —— *maculatus*, Johnston, p. 399, = *Gammaropsis maculatus* (Johnston).
106. *Megamæra semiserrata*, Bate, p. 401, = *Ceradocus semiserratus* (Bate).
107. —— *longimanus* (Leach, MS.), p. 403, = *Mæra othonis* (H. Milne-Edwards), ♂.
108. —— *othonis* (H. Milne-Edwards), p. 405, = *Mæra othonis* (H. Milne-Edwards), ♀.
109. —— *Alderi*, Bate, p. 407, = *Melita obtusata* (Montagu), ♀.
110. —— *brevicaudata*, Bate, p. 409, = *Elasmopus rapax*, Costa.
111. *Eiscladius longicaudatus*, Bate and Westwood, p. 412, = *Photis longicaudata* (Bate and Westwood).
112. *Amphithoe rubricata* (Montagu), p. 418.
113. —— *littoralis*, Bate, p. 422, = *Amphithoe rubricata* (Montagu), ♀.
114. —— *albonervulata*, Bate and Westwood (nec Kröyer), p. 426. The specimen figured is in my collection. It is *Podoceropsis excavata*, Bate, ♀. The long setæ of the antennæ and the first gnathopods as drawn agree with *Podoceropsis*, but the specimen shows nothing of the strong upturned third segment of metasome or the *Amphithoe*-like last uropod of Bate and Westwood's figure.
115. —— *gammaroides*, Bate, p. 427, = *Pleonexes gammaroides*, Bate, ♂.
116. *Sunamphithoe hamulus*, Bate, p. 430, = *Pleonexes gammaroides*, Bate, ♀.
117. —— *conformata*, Bate, p. 432, = *Sunamphithoe pelagica* (H. Milne-Edwards).
118. *Podocerus pulchellus* (Leach), p. 436, = *Bruzeliella* \* *falcata* (Montagu), ♂.

\* I have examined the types of the genera *Podocerus* and *Jassa* in the British Museum, and fully confirm what Messrs. Stebbing and Walker have written upon them, that *Podocerus variegatus*, Leach (as interpreted by Milne-Edwards's figure in 'Règne Animal de Cuvier,' but not *P. variegatus*, B. & W.), is *Platophium Darwini*, and that *Jassa pelagica*, Leach (but not of B. & W.), is the same as *Podocerus capillatus*, B. & W. Bruzelius long ago restricted the use of *Jassa* to the species last mentioned, which has only a rudimentary appendage (scarcely visible with a high power) to the antennules. This use must be maintained, and not *Parajassa* as suggested by Stebbing. I have therefore found it necessary to institute a genus *Bruzeliella*, with *B. falcata*, Montagu, as its type.

84 Canon A. M. Norman—*Revised Nomenclature of*

119. *Podocerus variegatus*, Bate and Westwood (nec Leach), p. 439, = *Bruzeliella pusilla* (Sars), ♀.
120. —— *capillatus*, Rathke, p. 442, = *Jassa pelagica*, Leach (nec B. & W.).
121. —— *falcatus* (Montagu), p. 445, = *Bruzeliella falcata* (Montagu), ♂ junior.
122. —— *pelagicus* (Bate and Westwood) (nec Leach), p. 447, = *Bruzeliella falcata* (Montagu), ♀.
123. —— *ocius*, Bate, p. 450, = *Bruzeliella ocia* (Bate).
124. *Cerapus abditus*, Templeton, p. 455, = *Erichthonius abditus* Templeton, ♂.
125. —— *diformis* (H. Milne-Edwards), p. 457, = *Erichthonius diformis*, H. Milne-Edwards, ♂.
126. *Derocthoe punctatus*, H. Milne-Edwards, p. 461, = *Erichthonius diformis*, H. Milne-Edwards, ♀.
127. *Siphonacetes typicus*, Bate and Westwood (nec Krüyer), p. 465, *Siphonacetes Whitei* (Gosse), ♂.
128. —— *Whitei*, Gosse, p. 467, = *Siphonacetes Whitei* (Gosse), ♀, = *C. Colletti*, Boeck.
129. —— *crassicornis*, Bate, p. 469, = *Cerapus crassicornis* (Bate).
130. *Nenia tuberculosa*, Bate, p. 472, = *Podoceropsis Sophieae*, Boeck, ♂.
131. —— *rimapalmata*, Bate, p. 474, = *Podoceropsis excavata* (Bate), ♂.
132. —— *excavata*, Bate, p. 476, = *Podoceropsis excavata* (Bate), ♀.
133. —— *undata*, Bate, p. 477, = *Podoceropsis Sophieae*, Boeck, ♀.
134. *Cyrtophium Darwinii*, Bate, p. 481, = *Platophium variegatum* (Leach)\*.
135. *Cratippus tenuipes*, Bate, p. 485, = *Colomastix pusilla*, Grube.
136. *Dryope irrorata*, Bate, p. 488, = *Unciola irrorata* (Bate), ♀.
137. —— *crenatipalma*, Bate, p. 490, = *Unciola irrorata* (Bate), ♂.
138. *Corophium longicorne*, Latreille, p. 493, = *Corophium volutator* (Pallas), ♂.
139. —— *Bonelli*, Bate and Westwood (nec Milne-Edwards), p. 497, = *Corophium affine*, Bruzelius, ♀.
140. —— *crassicornis*, Bruzelius, p. 499, ♂.
141. *Chelura terebrans*, Philippi, p. 503.

Vol. II.

142. *Listrigonus exulans*, Kröyer, p. 5, = *Hyperia galba* (Montagu), ♂ junior.

\* I think that, considering the inadequate description of the genus *Podocerus* and its erroneous use for nearly one hundred years, the name ought to be excluded from an altered use.

143. *Listrigonus Kinahani*, Bate, p. 8, = *Hyperia galba* (Montagu), ♂ mature.
144. *Hyperia galba* (Montagu), p. 12, ♀.
145. — *oblivia*, Bate und Westwood (nec Kröyer), p. 16, = *Euthemisto gracilipes* (Norman).
146. *Phronima sedentaria* (Forskaal), p. 23.
147. *Dulichia porrecta*, Bate, p. 31, ♂.
148. — *falcata*, Bate, p. 33, ♂.
149. *Proto pedata*, Abildgaard, p. 38, = *Phitisca marina*, Slabber, ♀.
150. — *Goodsirii*, Bate, p. 42, = *Phitisca marina*, Slabber, ♂.
151. *Protella phusma* (Montagu), p. 45, = *Pseudoprotella phasma* (Montagu).
152. *Caprella linearis* (Linné), p. 52.
153. — *lobata* (Müller), p. 57, = *Caprella linearis* (Linné), ♂.
154. — *acutifrons* (Latrille), p. 60.
155. — *hystrix* (Kröyer), Bate and Westwood, p. 63, = *Caprella septentrionalis*, Kröyer, var. *nodifera*, Mayer.
156. — *acanthifera*, Leach, p. 65.
157. — *tuberculata*, Guérin, p. 68.
158. — *aequilibra*, Say, p. 71.
159. — *typica*, Kröyer, p. 75, = *Periambus typicus* (Kröyer).
160. *Cyamus ceti*, Martens, p. 85, = *Cyamus mysticeti*, Lütken.
161. — *ovalis*, Roussel de Vauzème, p. 91, not British.
162. — *gracilis*, Roussel de Vauzème, p. 94, not British.
163. — *Thompsoni*, Gosse, p. 96, = *Platocyamus Thompsoni* (Gosse).

#### ISOPODA.

164. *Tanaid vittatus* (Rathke), p. 125, = *Tanaid Cavolini*, H. Milne-Edwards.
165. — *Dulongii* (? Audouin), p. 129.
166. *Leptochelia Edwardsii* (Kröyer), p. 134, = *Leptochelia Savignii* (Kröyer).
167. *Paratanais forcipatus*, Bate and Westwood (nec Lilljeborg), p. 138, = *Paratanais Batei*, G. O. Sars.
168. — *rigidus*, Bate and Westwood, p. 141, = *Leptognathia rigidula* (Bate and Westwood).
169. *Apseudes talpa* (Montagu), p. 148.
170. — *Latreillii*, H. Milne-Edwards, p. 153 (vide Norman, Ann. & Mag. Nut. Hist. ser. 7, vol. iii. p. 327).
171. *Anthura gracilis* (Montagu), p. 160, ♀.
172. *Paranthura Costana*, Bate and Westwood, p. 165, = *Paranthura nigropunctata* (Lucas), ♀.

173. *Anceus maxillaris* (Montagu), p. 187, = *Gnathia maxillaris* (Montagu).
174. — *Edwardii*, Bate, p. 201, = *Gnathia oxyuraa* (Lilljeborg).
175. — *Halidaii*, Bate and Westwood, p. 203, = *Gnathia formica* (Hesse).
176. *Bopyrus squillarum*, Latreille, p. 218.
177. *Gyge galatheae*, Bate and Westwood, p. 225, = *Gyge branchialis*, Cornalia and Panceri. The specimen which is described and illustrated at p. 225 was procured by me from *Upogebia stellata*, and not, as erroneously recorded by the authors, from a *Galathaea*.
178. — *hippolytes*, Kröyer, p. 230, = *Bopyroides hippolytes* (Kröyer).
179. *Phryxus abdominalis* (Kröyer), p. 234.
180. — *fusticaudatus*, Bate and Westwood, p. 238, = *Athelys paguri*, Rathke, young.
181. — *paguri*, Rathke, p. 240, = *Athelges paguri* (Rathke).
182. — *Hyndmanni*, Bate and Westwood, p. 243, = *Pseudione Hyndmanni* (Bate and Westwood).
183. — *longibranchiatus*, Bate and Westwood, p. 246, = *Pleurorypta longibranchiata* (Bate and Westwood).
184. — *galathae*, Bate and Westwood (nec Hesse), p. 249, = *Pseudione confusa* (Norman).
185. *Ione thoracica* (Montagu), p. 255.
186. *Cryptothiria pygmaea* (H. Rathke), p. 261, = *Liriopsis pygmaea* (H. Rathke).
187. — *balani* (Bate), p. 267, = *Cryptothir balani* (Bate).
188. *Ega bicarinata*, Leach, p. 278, = *Ega rosacea*, Risso. A doubtful British species (see Norman, Ann. & Mag. Nat. Hist. ser. 7, vol. xiv. 1904, p. 433).
189. — *tridens*, Leach, p. 281.
190. — *psora* (Pennant), p. 283.
191. — *monophtalma*, Johnston, p. 286.
192. *Rocinela danmoniensis*, Leach, p. 291.
193. *Cirolana Cranchii*, Leach, p. 296.
194. — *spinipes*, Bate and Westwood, p. 299, = *Cirolana borealis*, Lilljeborg.
195. *Conilera cylindracea* (Montagu), p. 304.
196. *Eurydice pulchra*, Leach, p. 310.
197. *Jæra albifrons* (Montagu), p. 317, = *Jæra marina* (Fabricius).
198. — *Nordmanni* (Rathke), p. 320.
199. *Munna Kröyeri*, Goodsir, p. 326, ♂.
200. — *Whiteana*, Bate and Westwood, p. 329, = *Munna Kröyeri*, Goodsir, ♀.

201. *Leptaspidea brevipes*, Bate and Westwood, p. 333. I think that this must be *Paramunna bilobata*, Sars, and should have precedence of that name.
202. *Janira maculosa*, Leach, p. 338.
203. *Asellus aquaticus* (Linné), p. 343.
204. *Limnoria lignorum* (Rathke), p. 351.
205. *Arcturus longicornis* (Sowerby), p. 363, = *Astacilla longicornis* (Sowerby), ♀.
206. — *intermedius* (H. Goodsir), p. 371, = *Astacilla intermedia* (H. Goodsir).
207. — *gracilis* (H. Goodsir), p. 373, = *Astacilla longicornis* (Sowerby), ♂.
208. *Idotea tricuspidata*, Desmarest, p. 379, = *Idotea balthica* (Pallas).
209. — *pelagica*, Leach, p. 384.
210. — *emarginata* (Fabricius), p. 386.
211. — *linearis* (Pennant), p. 388.
212. — *parallela*, Bate and Westwood, p. 391, = *Zenobiana prismatica* (Risso).
213. — *acuminata*, Leach, p. 394, = *Stenosoma acuminatum*, Leach.
214. — *appendiculata* (Risso), p. 396, = *Stenosoma lanciferum* (Leach).
215. *Sphaeroma serratum* (Fabricius), p. 405.
216. — *rujicauda*, Leach, p. 408.
217. — *Hookeri*, Leach, p. 410.
218. — *curtum*, Leach, p. 412, = *Cymodoce truncata* (Montagu), ♀.
219. — *Prideauxianum*, p. 415, = *Cymodoce truncata* (Montagu), ♀.
220. *Dynamene rubra*, Leach, p. 419, = *Næsa bidentata* (Adams), ♀.
221. — *viridis*, Leach, p. 421, = *Næsa bidentata* (Adams), ♀.
222. — *Montagni*, Leach, p. 423, = ?? young of *Næsa bidentata* (Adams), ♂.
223. *Cymodocea truncata*, Leach, p. 426, ♂, = *Cymodoce truncata*, Leach, ♂.
224. — *emarginata*, Leach, p. 428, = *O. truncata*, Leach, ♂ variety.
225. *Næsa bidentata* (Adams), p. 431, ♂.
226. *Campecopea hirsuta* (Montagu), p. 434, ♂.
227. — *Cranchii*, Leach, p. 436, = *Campecopea hirsuta* (Montagu), ♀.
228. *Ligia oceanica* (Liuné), p. 444.

229. *Philoscia muscorum* (Scopoli), p. 450.  
 230. —— *Couchii*, Kinahan, p. 452.  
 231. *Philouryria riparia* (Koch), p. 456, = *Trichoniscus pusillus*, Brandt.  
 232. —— *vividula* (Koch), p. 458, = *Trichoniscus vividus* (Koch).  
 233. —— *rosea* (Koch), p. 460, = *Trichoniscus roseus* (Koch).  
 234. *Platyarthrus Hoffmannseggii*, Brandt, p. 464.  
 235. *Oniscus asellus*, Linné, p. 468.  
 236. —— *fossor*, Koch, p. 471, = *Oniscus asellus*, Linné.  
 237. *Porcellio scaber*, Latreille, p. 475.  
 238. —— *dilatatus*, Brandt, p. 478.  
 239. —— *pictus*, Brandt, p. 480.  
 240. —— *laevis*, Latreille, p. 483.  
 241. —— *armadilloides*, Lereboullet, p. 485, = *Cylisticus convexus* (De Geer).  
 242. —— *pruinosus*, Brandt, p. 487, = *Metoponorthus pruinosus* (Brandt).  
 243. —— *cingendus*, Kinahan, p. 489, = *Metoponorthus cingendus* (Kinahan).  
 244. *Armadillo vulgaris*, Latreille, p. 492, = *Armadillidium vulgare* (Latreille).

## APPENDIX.

245. *Orchestia brevidigitata*, Bate and Westwood, p. 498, = *Orchestia littorea* (Montagu), ♂ young.  
 246. *Montaguea clypeata*, Bate and Westwood (nec Kröyer), p. 499, = *Metopa abscisa*, Normau.  
 247. —— *norvegica*, Bate and Westwood (nec Lilljeborg), p. 500, = *Metopa Alderi* (Bate), ♂.  
 248. *Opis leptochela*, Bate and Westwood, p. 501, = *Euonyx chelatus*, Norman.  
 249. —— *quadrifimana*, Bate and Westwood, p. 503, = *Normanion quadrifimanus* (Bate and Westwood).  
 250. *Ampelisca levigata*, Bate and Westwood (nec Lilljeborg), p. 504, = *Ampelisca tenuicornis*, Lilljeborg.  
 251. *Haploops tubicola*, Lilljeborg, p. 505.  
 252. *Monoculodes longimanus*, Bate and Westwood, p. 507, = *Peri-oculodes longimanus* (Bate and Westwood).  
 253. *Kroyera brevicarpa*, Bate and Westwood, p. 508, = *Synchelidium haplochiles* (Grube).  
 254. *Lepidepecreum carinatum*, Bate and Westwood, p. 509, = *Lepidepecreum longicorne* (Bate and Westwood), ♀.  
 255. *Nicripe tumida*, Bruzelius, p. 511.

256. *Cheirocratus mantis*, Norman, p. 513, = *Chirocratus assimilis* (Lilljeborg), ♂.  
257. *Megamara multidentata*, Bate and Westwood, p. 515, = *Mera Batei*, Norman.  
258. *Unciola leucomes*, Bate and Westwood (nec Kröyer), p. 517, = *Unciola planipes*, Norman.  
259. *Hyperoche tauriformis*, Bate and Westwood, p. 519, = *Hyperoche tauriformis* (Bate and Westwood).  
260. —— *prehensilis*, Bate and Westwood, p. 520, = ? *Hyperia tauriformis*, Bate and Westwood, young (*vide* Norman, Ann. & Mag. Nat. Hist. ser. 7, vol. v. 1900, p. 130).  
261. —— *cyanæ* (Sabine), p. 521, = ??  
262. *Themisto crassicornis*, Kröyer, p. 522, = *Euthemisto libellula* (Mandt).  
263. *Vibiliæ borealis*, Bate and Westwood, p. 524.  
264. *Tessarops hastata*, Norman, p. 530, = *Tiron acanthurus*, Lilljeborg.  
265. *Eriopis elongatus*, Bruzelius, p. 530, = *Eriopis elongata* (Bruzelius).  
266. *Mara Loveni*, Bruzelius, p. 530.  
267. —— *Batei*, Norman, p. 530.  
268. *Helleria coalita*, Norman, p. 530, = *Guernea coalita* (Norman).  
269. *Microprotopus maculatus*, Norman, p. 530.  
270. *Cirolana truncata*, Norman, p. 530, = *Eurydice truncata* (Norman).  
271. *Anilocra mediterranea*, Leach, p. 530, = *Anilocra physodes* (Linnae).

## II.

Species described by Bate and Westwood in accordance with present Nomenclature, and arranged after the Monographs of Professor G. O. Sars \*.

### I. AMPHIPODA.

#### Tribe I. HYPERIDEA.

*Hyperia galba* (Montagu), 142 and 143 ♂, 144 ♀.

— *cyanæ*, Sabine, 261 ?

*Hyperoche tauriformis* (B. & W.), 259 (and ? 260).

*Euthemisto libellula* (Mandt), 262.

— *gracilipes* (Norman), 145.

\* The numbers appended to the names are those prefixed to the species in the preceding list of Bate and Westwood's species.

*Phronima sedentaria* (Forskaal), 140.*Fibilia borealis*, B. & W., 263.

## Tribe II. GAMMARIDEA.

*Talitrus locusta* (Pallas), 1.*Orchestes littoreus* (Montagu), 2, 245.— *mediterranea* (Costa), 3.*Orchestoidea Deshayecii* (Audouin), 4.*Hyale Nilssonii* (Rathke), 5.— *Lubbockiana* (Bate), 6 ♂, 7 ♀.*Normanion quadrimanus* (B. & W.), 249.*Acidostoma obesum* (Bate), 19.*Lysianax septentrionalis*, Della Valle, 18 and part of 16.*Ambasia Danielsseni*, Boeck, 15?*Perrierella Audouiniana* (Bate), 14.*Callisoma Hopei*, Costa, 26.*Hippomedon denticulatus*, Bate, 20.*Orchomene humilis* (Costa), 18 and part of 16, and ? 22 junior.*Tryphosites longipes* (Bate), 24 ♀, 25 ♂.*Haplonyx cicada* (Fabricius), 21.*Menigrates obtusifrons*, Boeck, 23.*Lepidepecreum longicorne* (Bate), 17 ♂, 254 ♀.*Euonyx chelatus*, Norman, 248.*Bathypercia Guilliamsonia* (Bate), 81 ♀.— *Robertsoni*, Bate, 82 ♂.— *pelagica*, Bate, 83 ♂.*Haustorius arenarius* (Slabber), 42.*Urothoe marinus*, Bate, 44 ♀, 43 ♂.— *brevicornis*, Bate, 45 ♀.— *elegans*, Bate, 46 ♂.*Phoxocephalus Holbölli* (Kröyer), 30.— *simplex* (Bate), 29.*Harpinia neglecta*, G. O. Sars, 31.*Ampelisca typica* (Bate), 27.— *brevicornis* (Costa), 28.— *tenuicornis*, Lilljeborg, 250.*Haploops tubicola*, Lilljeborg, 251.*Amphilochus manudens*, Bate, 40.*Stenothoe marina* (Bate), 9.— *monoculoides* (Montagu), 8.

- Metopa Alderi* (Bate), 10 ♀, 247 ♂.  
 —— *norvegica* (Lilljeborg), 11.  
 —— *alscisa*, Norman, 246.  
*Cressa dubia* (Bate), 12.  
*Leucolhoe spinicarpa* (Müller), 70.  
 —— ? *farina* (Savigny), 71.  
*Monoculodes carinatus*, Bate, 36, 37.  
*Periocolodes longimanus* (B. & W.), 252.  
*Pontocrates arenarius* (Bate), 38.  
 —— *altamarinus* (Bate), 39.  
*Synchelidium haplochelis* (Grube), 253.  
*Halimedon parvimanus* (B. & W.), 33, 34, 35.  
*Parapleustes latipes* (M. Sars), 66, 67.  
*Epimeria cornigera* (Fabricius), 56.  
*Iphimedia obesa*, Rathke, 52.  
 —— *eblanae*, Bate, 53.  
*Pereionotus testudo* (Montagu), 55.  
*Odius carinatus* (Bate), 54.  
*Laphystius sturionis*, Kröyer, 41.  
*Tiron acanthurus*, Lilljeborg, 264.  
*Nicippe tumida*, Bruzelius, 255.  
*Eusirus longipes*, Boeck, 69.  
*Apherusa bispinosa* (Bate), 62.  
 —— *cirrus*, Bate, 63 ♂.  
 —— *Jurini* (H. M.-Edw.) ?, 72.  
*Calliopius laviusculus* (Kröyer), 65, 68 ♂.  
*Paratylus Swammerdamii* (H. M.-Edw.), 60.  
 —— *vedlomensis* (B. & W.), 59.  
*Dexamine spinosa* (Montagu), 57.  
 —— *thea*, Boeck, 59.  
*Tritata gibbosa* (Bate), 61 ♀.  
*Guernea coalita* (Norman), 268.  
*Amathilla homari* (Fabricius), 32 young, 97.  
*Gammarus marinus*, Leach, 98.  
 —— *campylops*, Leach, 99.  
 —— *locusta* (Linnaeus), 100.  
 —— *pulex*, De Geer, 103.  
 —— *Edwardsii*, Bate ?, 102.  
 —— " *dubius*, Johnston," 104.  
*Melita palmata* (Montagu), 90.  
 —— *obtusata* (Montagu), 91 and 92 ♂, 109 ♀.

- Melita glauiosa*, Bate, 93.  
*Eriopisa elongata* (Bruzelius), 265.  
*Niphargus subterraneus* (Leach), 84.  
 —— *fontanus*, Bate, 85.  
 —— *Kochianus*, Bate, 86.  
*Crangonyx subterraneus*, Bate, 87.  
*Mura othonis* (H. M.-Edw.), 108 ♀, 107 ♂.  
 —— *Loveni* (Bruzelius), 266.  
 —— *Batei*, Norman, 101 ?, 257, 267.  
 —— *grossimana* (Montagu), 94.  
*Ceradocus semiserratus* (Bate), 106.  
*Elasmopus rapax*, Costa, 110.  
*Gammarella brevicaudata*, 64 ♀, 88 ♂, 89 ♀.  
*Cheirocratus Sundevallii* (Rathke), 48 ♂, 80 ♀.  
 —— *assimilis* (Lilljeborg), 256 ♂.  
*Lilljeborgia pallida*, Bate, 47 ♀.  
 —— *Kinahani*, Bate, 50.  
" *Phaedra antiqua*, Bate," 49.  
*Isaea Montagui*, H. M.-Edw., 51.  
*Microdeutopus anomalus* (Rathke) ?, 77 ♀.  
 —— *dannoniensis* (Bate), 75 ♂.  
*Aoru gracilis*, Bate, 73.  
*Stimpsonella chelifera* (Bate), 74.  
*Lembos Websterii*, Bate, 76.  
*Coremapus versiculatus* (Bate), 78.  
*Leptocheirus pilosus*, Zaddach, 79.  
*Gammaropsis maculatus* (Johnston), 95 ♂, 96 ♀, 105.  
*Microprotopus maculatus*, Norman, 269.  
*Photis longicaudata* (Bate), 111.  
*Podoceropsis Sophie*, Boeck, 130 ♂, 133 ♀.  
 —— *excavata* (Bate), 131 ♂, 132 ♀, 114.  
*Amphithoe rubricatu* (Montagu), 112 ♂, 113 ♀.  
*Pleonexes gammaroides*, Bate, 115 ♂, 116 ♀.  
*Sunamphithoe pelagica* (H. M.-Edw.), 117.  
*Bruzeliella fulcata* (Montagu), 121 ♂ junior, 118 ♂, 122 ♀.  
 —— *pusilla* (G. O. Sars), 119 ♀.  
 —— *ocia* (Bate), 123.  
*Jassa pelagica*, Leach, 120.  
*Erichthonius alditus* (Templeton), 124 ♂.  
 —— *diformis* (H. M.-Edw.), 125 ♂, 126 ♀.  
*Cerapus crassicornis* (Bate), 129.

- Siphonæcetes Whitei* (Gosse), 127 ♂, 128 ♀.  
*Corophium volutator* (Pallas), 138 ♂.  
 —— *affine*, Bruzelius, 139 ♀.  
 —— *crassicornis*, Bruzelius, 140 ♂.  
*Unciola irrorata* (Bate), 136 ♀, 137 ♂.  
 —— *planipes*, Norman, 259.  
*Chelura terebrans*, Philippi, 141.  
*Platophium variegatum* (Leach), 134.  
*Dulichia porrecta*, Bate, 147 ♂.  
 —— *falcata*, Bate, 148 ♂.  
*Colomastix pusilla*, Grube, 135.

## Tribe III. CAFRILLIDEA.

- Phthisica marina*, Slabber, 149 ♀, 150 ♂.  
*Pseudoprotella phasma* (Montagu), 151.  
*Periambus typicus* (Kröyer), 159.  
*Caprella linearis* (Linné), 152 ♀, 153 ♂.  
 —— *septentrionalis*, Kröyer, var. *nudifera*, Mayer, 155.  
 —— *aequilibra*, Say, 158.  
 —— *acanthifera*, Leach, 156.  
 —— *acutifrons* (Latreille), 154.  
 —— *tuberculata*, Guérin, 157.  
*Cyamus mysticeti*, Lütken, 160.  
 —— ? *ovalis*, Roussel de Vauzème, 161.  
 —— ? *gracilis*, Roussel de Vauzème, 162.  
*Platycyamus Thompsoni* (Gosse), 163.

## II. ISOPODA.

## Tribe I. CHELIFERA.

- Apseudes talpa* (Montagu), 169.  
 —— *Latreillii* (H. M.-Edw.), 170.  
*Tanaïs Carolini*, H. M.-Edw., 164.  
 —— *Dulongji* (? Audouin), 165.  
*Leptochelia Savignii* (Kröyer), 166.  
*Paratanais Batei*, G. O. Sars, 167.  
*Leptognathia rigida* (B. & W.), 168.

## Tribe II. FLABELLIFERA.

- Anthura gracilis* (Montagu), 171.  
*Paranthura nigropunctata* (Lucas), 172.

94        *Crustacea described by Bate and Westwood.*

- Gnathia maxillaris* (Montagu), 173.  
— *oxyurata* (Lilljeborg), 174.  
— *formica* (Hesse), 175.  
*Anilocra physodes* (Linné), 271.  
*Aega tridens*, Leach, 189.  
— *psora* (Pennant), 190.  
— *monophthalma*, Johnston, 191.  
— ? *rosacea*, Risso, 188.  
*Rocinela ammoniensis*, Leach, 192.  
*Cirolana Cranchii*, Leach, 193.  
— *borealis*, Lilljeborg, 194.  
*Conilera cylindracea*, Leach, 195.  
*Eurydice pulchra*, Leach, 196.  
— *truncata* (Norman), 270.  
*Limnoria lignorum* (Rathke), 204.  
*Sphaeroma serratum* (Fabricius), 215.  
— *rugicauda*, Leach, 216.  
— *Hookeri*, Leach, 217.  
*Cymoloea truncata*, Leach, 218 and 219 ♀, 223 and 224 ♂.  
*Nasa bidentata* (Adams), 220 and 221 ♀, 225 ♂.  
— *Montaqui*, Leach ?, 222.  
*Campecopea hirsuta* (Montagu), 226 ♂, 227 ♀.

Tribe III. VALVIFERA.

- Astarilla longicornis* (Sowerby), 205 ♀, 207 ♂.  
— *intermedia* (Goodsir), 206.  
*Idotea balthica* (Pallas), 208.  
— *pelagica*, Leach, 209.  
— *emarginata* (Fabricius), 210.  
— *linearis* (Pennant), 211.  
*Zenobiana prismatica* (Risso), 212.  
*Stenosoma acuminatum*, Leach, 213.  
— *lanciferum* (Leach), 214.

Tribe IV. ASELLOTA.

- Asellus aquaticus* (Linné), 203.  
*Janira maculosa*, Leach, 202.  
*Jara marina* (Fabricius), 197.  
— *Nordmanni* (Rathke), 198.  
*Munna Kröyeri*, Goodsir, 199 ♂, 200 ♀.  
*Leptaspidea brevipes*, B. & W., 201.

## Tribo V. ONISCOIDEA.

- igia oceanica* (Linné), 228.  
*richoniscus pusillus*, Brandt, 231.  
 — *vividus* (Koch), 232.  
 — *roseus* (Koch), 233.  
*Oniscus asellus*, Linné, 235, 236.  
*Philoscia muscorum* (Scopoli), 229.  
 — *Couchii*, Kinahan, 230.  
*Platyarthrus Hoffmannseggii*, Brandt, 234.  
*Porcillio scaber*, Latreille, 237.  
 — *dilatatus*, Brandt, 238.  
 — *pictus*, Brandt, 239.  
 — *levis*, Latreille, 240.  
*Metoponorthus pruiniosus* (Brandt), 242.  
 — *cinctus* (Kinahan), 243.  
*Cylisticus conicus* (De Geer), 241.  
*Armadillidium vulgare* (Latreille), 244.

## Tribe VI. EPICARIDA.

- Bopyrus squillarum*, Latreille, 176.  
*Pseudione Hyndmanni* (B. & W.), 182.  
 — *confusa* (Norman), 184.  
*Pleurocrypta longibranchiata* (B. & W.), 183.  
*Athalyes paguri* (Rathke), 181 and 180, the young.  
*Phryxus abdominalis* (Kröyer), 179.  
*Ione thoracica* (Montagu), 185.  
*Liriopsis pygmaea* (Rathke), 186.  
*Cryptothir balani* (Bate), 187.
- 

IX.—*Kerunia, a Symbiosis of a Hydractinian with a Cephalopod.* By DR. FRANCIS BARON NOPCSA, Member of the Hungarian Geological Society.

[Plate III.]

DURING a recent visit to Qasr-el-Sagha, in the Fayum district of Egypt, I succeeded in obtaining a large number of specimens belonging to the Eocene genus *Kerunia* uniting characters which have hitherto not been properly described.