

*5
R Fishes*

Canceled

Division of Fishes,
U. S. National Museum

BULLETIN

OF THE

UNITED STATES FISH COMMISSION.

VOL. XI.

FOR

1891.

MARSHALL McDONALD, Commissioner.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1893.

10.—ON THE CLASSIFICATION OF THE MYXOSPORIDIA, A GROUP OF PROTOZOAN PARASITES INFESTING FISHES.

BY R. R. GURLEY, M. D.,
Assistant, U. S. Fish Commission.

Up to the present time very little attention has been given to the diseases of fishes and to their parasites from the standpoint of the effect produced upon the host; yet there can be no doubt that a knowledge of such diseases would be of great practical value. Anyone who considers the proportions that fish epidemics sometimes attain will hardly be inclined to question the utility of searching investigation in this direction. Thus, to take a single instance, Prof. Forbes states* that in the epidemic of 1884 in Lake Mendota it was estimated that fully 300 tons had died. On August 7 the Madison Transcript reported that 200 tons of fish had been hauled away by the city authorities during the four weeks preceding and that the fishes were still dying. Epidemics of similar extent have been reported in Europe, for several of which (that of the barbel certainly, and that of the crayfish probably) the *Myxosporidia* are responsible.

The important results in the way of prevention of epidemics among domesticated animals and cultivated plants, obtained as the result of scientific investigation, afford ground for the hope that similar results may be obtained here. Obviously the first step in work of this kind is the collection of facts, especially those bearing upon the parasite, its nature, life history, intermediate hosts, enemies, and its connection (whether causal or otherwise) with diseases or other morbid processes in its host. Such data are a necessary preliminary to preventive or curative measures.

The present paper and a more extended one now in preparation are intended as contributions to the objects indicated. In the latter paper the practical bearings of the subject will be fully discussed, and all the data as to epidemics of myxosporidiosis will be given. At present it is desired mainly to discuss the classification of the subclass *Myxosporidia* Bütschli and to record such genera and species as a study of the literature and of such material as was available has led me to recognize. These forms will all be fully described and figured in the second paper. The present only includes such true *Myxosporidia* as have received or appear entitled to receive binomial names; and only such synonymy as is needed for their identification.

* Bull. U. S. Fish Com. 1888, VIII, p. 482.

CLASSIFICATION OF THE SUBCLASS.

The only classification ever proposed is that of Thélohan.* This author enunciates three taxonomic principles:

1. The habitat furnishes no sound basis for specific distinctions.†
2. The myxosporidium affords no taxonomic criteria.
3. The spores alone offer (at least in the present state of our knowledge) characters suitable to serve as a basis for classification. He says:

By noting the differences of form and size of these elements, the number of their polar capsules, by taking account of the presence or absence of a vacuole in the plasma, of their number in the [pan]sporoblasts‡, one can, I believe, succeed in obtaining elements sufficient for an attempt of this kind.

Thélohan further states that he regards the classification only as a provisional one, and that it is the result of a desire to obviate the great confusion arising from the habit of designating forms by their habitats.

The following is Thélohan's primary classification:

MYXOSPORIDIANS.

Spores	{	Pyriform; capsule 1, at pointed extremity; vacuole 1, aniodinophile, at large extremity.	} I. <i>Glugeidians</i> .
		Form variable	{
		No vacuole; capsules 2 or 4.	{ Capsules 2 II. <i>Myxidians</i> . [*] Capsules 4 III. <i>Chloromyxans</i> .
		Vacuole 1, iodophilic.	Capsules 1-2 IV. <i>Myxobolans</i> .

The three principles mentioned, which form the basis of Thélohan's classification, are unquestionably sound as far as they go, but to obtain a satisfactory arrangement it will sometimes be necessary to employ additional characters and to arrange them in a different order of rank. Especially will it be necessary in the *Phænocystes* to refer constantly to the symmetry and topography of the spore.

Further, while there can be no question as to the propriety of drawing a sharp line between the "Glugeidians" of the above table and the remaining *Myxosporidia*,

*Bull. Soc. philomat. Paris, 1892, iv, pp. 165-178.

† While as a general principle this is beyond question, indications are not wanting to show that in some cases the seat and, to a somewhat less extent, the host, bear some relations to generic lines. One of the most significant facts of organal and zoölogical distribution is the following:

Forms.	Total number of species.	Gall and urinary bladders, bile ducts and urinary tubules.	Remaining organs.
Non-vacuolous	9	8	1
Vacuolous	Very many . . .	<i>Ad plur</i> 2 (3 ?)	All but 2 (or 3).

‡ *Pansporoblast*: The transparent plasma-sphere formed by the condensation of a portion of the plasma around one of the numerous nuclei of the endoplasm of the myxosporidium; in distinction from the sporoblasts which result from the segmentation of the pansporoblast.

the division certainly can not rest upon such a comparatively unimportant character as the shape of the spore. I have regarded this division as of ordinal value and define the two orders thus:

I. *Cryptocystes* ord. nov. *Myxosporidia* in which the pansporoblast produces many (at the fewest 8) spores; the last minute, without distinct symmetry, with a single capsule; type (and only) family, *Glugeidæ* fam. nov.

Etymology: *κρυπτός*, concealed; *κύστις*, capsule.

II. *Phænocystes* ord. nov. *Myxosporidia* in which the pansporoblast produces few (at the most 2) spores;* the last relatively large, with distinct symmetry and 2 or more capsules;† type family, *Myxobolidæ* fam. nov.

Etymology: *φαίνω*, I appear; *κύστις*, capsule.

I. CRYPTOCYSTES ord. nov.

GLUGEIDÆ fam. nov.

Glugeidées Thélohan, 1892, Bull. Soc. philomat. Paris, iv, pp. 173-174.

Definition (provisional as regards negative characters): *Cryptocystes* destitute of a bivalve shell; with the capsule at the anterior extremity and with an aniodinophile vacuole; type genus, *Glugea* Thélohan.

This family contains three genera,‡ whose relations are shown in the following table:

Myxosporidium.	Spores formed in pansporoblast.	Pansporoblast membrane.	Genera.
Present.....	Inconstant, numerous.....	Not subpersistent.....	<i>Glugea</i> Thélohan.
Absent.....	Inconstant, numerous.....	Subpersistent.....	<i>Pleistophora</i> gen. nov.
Absent.....	Constant, 8.....	Subpersistent.....	<i>Thélohania</i> Henneguy.

GLUGEA Thélohan, 1891.

Compt. Rend. hebdom. Soc. Biol. Paris, III, p. 29; *ib.* Thélohan, 1892, Bull. Soc. philomat. Paris, iv, p. 174.

Definition: *Glugeidæ* possessing a myxosporidium, and in which the pansporoblast produces an inconstant but large number (always more than 8) of spores; pansporoblast membrane not subpersistent; type, *G. microspora* Thél. (synonym for *G. anomala* Moniez).

Glugea anomala Moniez, 1887.

Nosema anomala, Compt. Rend. Acad. Sci. Paris, CIV, p. 1312; *Glugea microspora*, Thélohan, 1891, Compt. Rend. hebdom. Soc. Biol. Paris, III, p. 29.

Glugea destruens Thélohan, 1892.

Bull. Soc. philomat. Paris, iv, pp. 165, 174 (footnote).

* Three asserted in one species (Leydig, Müller's Archiv., 1851, p. 229).

† Except (at most) two *Myxobolus* species (one of them perhaps inconstantly), which have suffered reduction to one.

‡ Thélohan recognizes only 2 genera, the distinctions between which are mainly based upon the three characters noted. If (as both he and I believe) these characters are sufficient to determine genera at all, a third genus must be recognized, as *Pleistophora typicalis* sp. nov. could not (as the above table shows) well be forced into either of the existing ones.

PLEISTOPHORA * gen. nov.

Definition (provisional as regards negative characters): *Glugeidae* destitute of a myxosporidium and in which the pansporoblast produces an inconstant but large number (always more than 8) of spores; pansporoblast membrane subpersistent (as a polysporophorous vesicle); type (and only) species, *P. typicalis* sp. nov.

Pleistophora typicalis sp. nov.

(Corpuscles of *Cottus scorpio* Thélohan, 1890, Annal. de Microgr. Paris, II, pp. 203, 212; *ib.* Thélohan, 1891, Compt. Rend. hebdom. Soc. Biol. Paris, III, pp. 27-8; *ib.* Henneguy and Thélohan, 1892, Annal. de Microgr., IV, pp. 618-619, 622, 631, 636.)

Pansporoblast: Spherical, diameter 15 to 18 μ .

Spore: Ovoid; length 3 μ ; breadth 1.5 to 2.0 μ ; chromatophile granules *ad plur.* 4.

Habitat: Interior of fibrillæ of muscles of *Cottus scorpio*; diseased mass forming white streaks 5 to 6 by 3 mm.; not leading to muscle degeneration.

THÉLOHANIA Henneguy, 1892.

In Thélohan, Bull. Soc. philomat. Paris, IV, p. 174 (footnote); *ib.* Henneguy and Thélohan, 1892, Annal. de Microgr., IV, p. 639.

Definition (provisional as regards negative characters): *Glugeidae* destitute of a myxosporidium and in which the pansporoblast produces constantly 8 spores; pansporoblast membrane subpersistent (as an octosporophorous vesicle); type, *T. giardi* Henneguy.†

Thélohania contejeani Henneguy, 1892.

In Thélohan, Bull. Soc. philomat. Paris, IV, p. 174 (footnote).

Thélohania octospora Henneguy, 1892.

In Thélohan, Bull. Soc. philomat. Paris, IV, p. 174 (footnote).

Thélohania giardi Henneguy, 1892.

In Thélohan, Bull. Soc. philomat. Paris, IV, p. 174 (footnote).

Thélohania macrocystis sp. nov.

(Sarcosporidian of *Palæmonetes varians* Garbini, 1891, Atti Reale Accad. Lincei Roma, VII, Sem. 1, pp. 151-152, with figs.; myxosporidian of *ibid.* Thélohan and Henneguy, 1892, Compt. Rend. hebdom. Soc. Biol. Paris, IV, p. 586).

Sporophorous vesicle (subpersistent pansporoblast) elongate-fusiform.

Habitat: Muscles of *Palæmonetes varians* from the Mincio, near Verona.

II. PHÆNOCYSTES ord. nov.

It is in the classification of this order that the criteria furnished by Thélohan most need to be supplemented by considerations drawn from the *symmetry* of the spore. Considering the taxonomic importance of symmetry throughout the animal kingdom, it is strange that no attention has yet been paid to it in the *Myxosporidia*. But a little study of it shows that, whereas in all fusiform spores the pointed extremities have heretofore been loosely lumped together as "ends," those of *Myxidium lieberkühni* Bütschli are not ends (*sens strict.*; = anterior and posterior), but sides; for the

* Etymology: *πλειστος*, very many; *φέρειν*, to carry.

† I propose *T. giardi* Henneguy as the type of the genus.

spore is symmetrical on either (right or left) side of the vertical plane,* but it is asymmetric on either (anterior and posterior) side of the transverse plane. On the other hand, if, as seems probable, the generic reference of *C.?* *diploxyis* sp. nov. be correct, then "ends" in *Cystodiscus* are ends, properly speaking. It is needless to emphasize the taxonomic import of these results, for we are thus enabled to orient the spore and the results of such orientation may be summed up as follows:

1. Within this order the most important characters are the *position* and *grouping* of the capsules. Compared to this the mere *number* of these bodies is a character of minor importance; for not only has *Myxobolus* 1 or 2 and *Cystodiscus* 2 or 4, but the number even varies in the same species, *Myxidium lieberkühni* Bütschli having 2 or 4.† But whether 1 to 2 or 2 to 4, the topographic relations are never varied. Thus in *Myxobolus* they are always in one group at the anterior end; in *Cystodiscus* in two groups, *anterior* and *posterior*; and in *Myxidium* in 2 groups, *right* and *left*.

Similar results are obtained with relation to the position of the valves, or, in other words, to the orientation of their plane of junction.

The following table shows the relations of these points to generic lines:

Comparison of generic characters in the *Phanocystes*.

[× = present; 0 = absent; () = less usual; — = condition not known.]

	Symmetry.		Capsules.				Shell.			Vacuole.	Tail.
	Antero-posterior.	Bilateral; perfect.	Number.	In one group (at the anterior end.)	In two groups.		Bi-valve.	Inclination of plane of junction of valves to longitudinal plane.			
					At the (anterior and posterior) ends.	In the (right and left) wings.		0°.	90°.		
<i>Cystodiscus</i> Lutz.....	×	×	2 (or 4)	×	×	×	—	0	
<i>Myxobolus</i> Bütschli <i>sens. strict</i> ..	0	×	2 (or 1)	×	×	×	0	
<i>Henneguya</i> Thélohan.....	0	×	2	×	×	×	×	
<i>Chloromyxum</i> Mingazzini.....	0	×	4	×	×	—	(?)	0	
<i>Mixosoma</i> Thélohan.....	0	×	2	×	— ²	—	—	0	
<i>Sphaerospora</i> Thélohan.....	0	×	2	×	×	—	—	0	
<i>Ceratomyxa</i> Thélohan.....	0	(¹)	2	×	×	×	0	
<i>Myxidium</i> Bütschli.....	0	×	2 (or 4)	×	0	0	

¹ Imperfect. Shell and capsules symmetrical; sporoplasm unilateral.

² From analogy and general similarity of appearance, this genus can hardly be other than bivalve.

*The three planes to which symmetry may be referred may be thus defined:

Vertical plane: Median, longitudinal and intercapsular in position.

Transverse plane: Vertical, transverse and (in *Myxobolus*) postcapsular in position.

Longitudinal plane: Horizontal, longitudinal and percapsular in position.

† Balbiani, 1883, Journ. de Microgr. Paris, VII, p. 274, fig. 64 g.

From this table we may conclude that—

1. *Cystodiscus* Lutz is certainly entitled to separate family rank.
2. *Henneguya* Thélohan agrees with *Myxobolus* in every respect but one, the presence of a tail.
3. Thélohan's groups, "*Myxidiées*" and "*Chloromyxées*" must undergo rearrangement (see table below); for clearly *Chloromyxum* Mingaz, *Mixosoma* Thél., and *Sphærospora* Thél. form a compact group, with which *Myxidium* has no character of consequence in common except the absence of a vacuole.
4. *Sphærospora* and *Mixosoma* do not differ at all in the characters given (the distinction between these unispecific genera resting solely upon the shape of the spore), and the two taken together present only a single character in contrast to *Chloromyxum*, viz, the number of the capsules.
5. *Ceratomyxa* agrees sufficiently closely with *Chloromyxum* to permit its reference to the *Chloromyxidæ*.
6. *Myxidium* must form the type of a separate family.

The following table shows the relations of Thélohan's classification to the one now proposed:

Thélohan's classification.			Proposed classification.		
	GROUP.		GENUS.	FAMILY.	CHARACTERS.
No vacuole, 2 or 4 capsules.	2 capsules. (II. Myxidians.)	Fusiform, 1 capsule at each extremity. <i>Myxidium</i> Bütschli.	<i>Myxidium</i> Bütschli.	<i>Myxidiidæ</i>	Bilateral but not antero-posterior symmetry; capsules in 2 groups right and left; no bivalve shell; no vacuole.
		Elongated; shell, formed of 2 hollow cone valves soldered along their bases. <i>Ceratomyxa</i> Thél.	<i>Ceratomyxa</i> Thél ..		
	Spores.	Flattened-ovoid, more or less elongate. <i>Mixosoma</i> Thél. Spherical. <i>Sphærospora</i> Thél.	<i>Chloromyxum</i> Mingaz, et sub-gen. <i>Sphærospora</i> .	<i>Chloromyxidæ</i> .	
	4 capsules. (III. Chloromyxans.)	<i>Chloromyxum</i> Mingaz.			
One iodophile vacuole; 1 or 2 capsules.	IV. Myxobolans.	Destitute of a tail; capsules 1 or 2. <i>Myxobolus</i> Bütschli.	<i>Myxobolus</i> Bütschli.	<i>Myxobolidæ</i>	Bilateral but not antero-posterior symmetry; capsules in 1 group (at the anterior end); a bivalve shell with the valve-junction plane parallel to the longitudinal plane, an iodophile vacuole.
		With a tail; capsules 2. <i>Henneguya</i> Thél.			
	Spore-shell.		<i>Cystodiscus</i> Lutz.	<i>Cystodiscidæ</i> ..	Bilateral and antero-posterior symmetry; capsules in 2 groups, anterior and posterior; a bivalveshell with the valve-junction plane perpendicular to the longitudinal plane; condition of sporoplasm unknown.

As a result of this analysis, the order *Phænocystes* may be divided into the following families:

CYSTODISCIDÆ fam. nov.

Definition: *Phænocystes* whose spores possess antero-posterior and bilateral symmetry; capsules in 2 groups, situated at the (anterior and posterior) ends; a bivalve shell, the plane of junction of whose valves is perpendicular to the longitudinal plane; condition of sporoplasm* unknown; type (and only) genus *Cystodiscus* Lutz.

CYSTODISCUS Lutz, 1889.

Centralbl. f. Bakt. u. Parasitenkde., v, p. 88.

Definition: Characters those of the family; type, *C. immersus* Lutz.

Cystodiscus immersus Lutz, 1889.

Centralbl. f. Bakt. u. Parasitenkde., v, pp. 84-88; figs. separately and subsequently.

Cystodiscus ? *diploxys* sp. nov.

(Psorosperms of *Pyralis viridana* Balbiani 1867, Journ. Anat. et Physiol. Paris, p. 335 (footnote), t. 12, f. 10-12.)

Spore: Parallel-sided fusiform; ends symmetrically double convex-curved pointed; plane of junction of valves coincident with the vertical plane; capsules 2 at each end, of equal size.

Habitat: *Tortrix viridana*.

MYXOBOLIDÆ fam. nov.

(*Myxosporidieæ*† Perugia, 1891, Boll. Scientif. Pavia, XIII, p. 23; "Myxobolées" Thélohan, 1892, Bull. Soc. philomat. Paris, IV, pp. 173, 176.)

Definition: *Phænocystes* whose spores are destitute of antero-posterior, but possess bilateral, symmetry;‡ capsules in 1 group at the anterior end; a bivalve shell, the plane of junction of whose valve is parallel to the longitudinal plane; an iodophile vacuole; type (and only) genus, *Myxobolus* Bütschli.

MYXOBOLUS Bütschli, 1882.

Bronn's Thier-Reich, I, t. 38, f. 6-10; *Myxosporidium* § Perugia, 1891, Boll. Scientif. Pavia, XIII, p. 23; *Myxobolus* et *Myxosporidium* Weltner, 1892, Sitzgsber. d. Gesellsch. Naturf. Freunde Berlin, pp. 34-35; *Myxobolus* et *Henneguya*|| Thélohan, 1892, Bull. Soc. philomat. Paris, IV, pp. 176, 177.

Definition: Characters those of the family; type *M. mülleri* Bütschli.

* *Sporoplasm.* Protoplasm of the spore.

† I propose *Myxosporidium* Perugia (synonym for *Myxobolus* Bütschli) as the type genus of the Fam. *Myxosporidieæ* Perugia.

‡ Except species which have suffered reduction of characters (*Myxobolus unicapsulatus* sp. nov., *M. piriformis* Thél.).

§ I propose *M. merlucii* Per. as the type species of this genus.

|| I propose *H. psorospermica* Thél. as the generic type. No valid generic distinction seems possible between the untailed and the tailed forms, for which latter Thélohan proposed *Henneguya*. Apart from the absence or presence of a tail (both of which conditions may, according to good observers, occur in the same species; cf. Weltner, Sitzgsber. Ges. Naturf. Freunde Berlin, 1892, pp. 28-36) the only character relied upon for their separation is the constancy of 2 capsules in the tailed forms, but this is also the typical number for *Myxobolus* and the presence of two exceptional species does not militate against the rule.

Myxobolus unicusulatus sp. nov.(Psorosperm of *Labeo niloticus* Müller, Müller's Archiv., 1841, p. 487, t. 16, f. 5 a-d.)

Spore: Of the form and size of *Chloromyxum dujardini* Thél.; capsule only 1, obliquely directed.

Habitat: *Labeo niloticus*, from the Nile.

Myxobolus piriformis Thélohan, 1892.

Bull. Soc. philomat. Paris, IV, p. 177.

Myxobolus inequalis sp. nov.(Psorosperms of *Pimelodus blochii* (Valenc.) Müller, Müller's Archiv., 1841, p. 487, t. 16, f. 6 a. b.)

Spore: Length 11 μ ; breadth, 7 μ ; capsules 2 of *unequal* size.

Habitat: On *Pimelodus clarias* Bloch (= *Silurus clarias* Valenc.), from Guiana and Surinam.

Myxobolus mugilis Perugia, 1891.*Myxosporidium mugilis*, Boll. Scientif. Pavia, XIII, pp. 23-4; *Myxobolus mugilis* Thélohan, 1892, Bull. Soc. philomat. Paris, IV, p. 166.**Myxobolus oviformis** Thélohan, 1892.

Bull. Soc. philomat. Paris, IV, p. 177.

Myxobolus mülleri Bütschli, 1882.

Bronn's Thier-Reich, I, t. 38, f. 6-10.

Myxobolus oblongus sp. nov.(Psorosperms of *Catostomus tuberculatus* Müller, Müller's Archiv., 1841, pp. 487-90, t. 16, f. 7-9.)

Cyst: Round or elliptic, 1 mm. or less.

Spore: Spatular or round-oblong; length, 14 to 17 μ ; greatest breadth and greatest thickness at about the junction of the anterior and second fourth of the length; breadth, 8.5 μ ; thickness, 5 to 6 μ ; width of ridge nearly equal to one-third of thickness; capsules, 2; nuclei 2, and perhaps* more; vacuole not seen.

Habitat: Subcutaneous on head of *Erimyzon sucetta* Lac. (= *Catostomus tuberculatus* Le Sueur).

Myxobolus ellipsoides Thélohan, 1892.

Bull. Soc. philomat. Paris, IV, p. 177.

Myxobolus bicostatus sp. nov.(Myxosporidian spore of branchiæ of *Tinca vulgaris* Bütschli, 1882; Bronn's Thier-Reich, I, t. 38, f. 19.)

Spore: Shell with 2 oblique ribs over the longitudinal axes of the capsules; capsular index 0.50. †

Habitat: Branchiæ of *Tinca tinca* L.

Myxobolus lintoni sp. nov.(Psorosperms of *Cyprinodon variegatus* Linton, 1891, Bull. U. S. Fish Com. for 1889, IX, pp. 99-102, t. 35, f. 1-16.)

Spore: Ovate; length, 13.9 μ ; breadth, 11 μ ; thickness about 8 μ ; shell, valves separating rapidly in sulphuric acid (cold, concentrated); capsules, 2, equal; filaments extruded under influence of sulphuric acid and of iodine water; nuclei *ad plur.* 4, 2 of which are the pericornual; ‡ vacuole present.

* The condition of the material being such as not to exclude the possibility of sporoplasmic degeneration, these results are not entirely decisive.

† *Capsular index.* The ratio of the length of the capsule to the antero-posterior diameter of the shell cavity.

‡ *Pericornual nuclei.* The 2 nuclei ("granules" "globules") situated at the antero-lateral angles (*cornua*) of the sporoplasm or on the posterior extremities of the capsules.

Habitat: Subcutaneous, on *Cyprinodon variegatus*, from the Atlantic at Woods Holl, Mass., August 20, 1889.

Myxobolus obesus, sp. nov.

(Psorosperm of the "Ablette," Balbiani, 1883, Journ de Microgr., VII, p. 203, fig. 43.)

Spore: Very broadly ovate.

Habitat: On *Alburnus alburnus* L.

Myxobolus cycloides, sp. nov.

(Psorosperms of *Cyprinus rutilus* (*pars*) Müller, Müller's Archiv., 1841, pp. 481, 486, t. 16, f. 4 d-g.)

Spore: Subcircular-ovate to broadly rounded-elliptic.

Habitat: Opercle and pseudobranchiæ of *Leuciscus rutilus* L., from German rivers, May and June.

Myxobolus spherialis sp. nov.

(Psorosperms of *Coregonus fera* Claparède, 1874, in Lunel's Hist. Nat. d. Poissons du bassin du Léman, Geneva, pp. 113-4.)

Spore: Untailed; perfectly spherical; diameter, 9μ ; containing a single spherical very strongly refringent "nucleus" [? vacuole].

Habitat: Cysts imbedded by thousands in the branchial mucosa of *Coregonus fera* Jur.

Myxobolus globosus, sp. nov.

Cyst: Elongate-elliptic or rod-shaped; length *ad max.*, 0.50 mm.

Spore: Globose; length, 7 or 8 μ ; breadth, 6 μ ; thickness, 5 μ ; ridge very wide, one-third of thickness; capsular index somewhat more than 0.50.

Habitat: Branchial lamellæ of *Erimyzon sucetta* Lac. (= *Catostomus tuberculatus* Le Sueur), from Kinston, N. C., Columbia, S. C., and from the tributaries of the Fox River, Miss. (collections U. S. National Museum).

Myxobolus transovalis, sp. nov.

Spore: Transversely elliptic; length, usually 6, occasionally 7 μ ; breadth, 8 μ ; valves separating in sulphuric acid; ridge narrow; capsules 2, equal; capsular index, 0.50; filaments extruded under the influence of glycerine and of sulphuric acid; nuclei, 2, rarely only 1, pericorncular nuclei apparently absent; vacuole probably present.

Habitat: In hollow of under surface of scales of *Phoxinus funduloides* from Four mile Run (tributary of Potomac River), Carlins, Va., June 29, 1892.

Myxobolus merlucii Perugia, 1891.

Myxosporidium merlucii, Boll. Scientif. Pavia, XIII, pp. 22, 24, f. 9-14.

Myxobolus perlatus, sp. nov.

(Psorosperms of *Acerina cernua* Balbiani, 1883, Journ. de Microgr. Paris, VII, pp. 201, 204, fig. 44.)

Spore: Wider than long.

Habitat: On *Gymnocephalus cernua* L.

Myxobolus?? zschokkei, sp. nov.*

(Psorosperms of *Coregonus fera* Zschokke, 1884, Archiv. de Biol. v, pp. 234-5, t. 10, f. 16.)

Cyst: Oval, pea to nut size; multiple (up to 30).

Spore: Body lenticular or oval, a little wider in front than behind; often bearing in front a blunt prolongation; posteriorly one distinguishes two "tails" (*queues*) 6 to 8 times longer than the body, attenuating posteriorly, curved and undulating; the number of two "tails" is constant; at the pole opposite to the "tails" are 2 oval, transparent anteriorly converging vesicles; one sometimes sees, however, an extremely fine canal extending from the posterior end of each vesicle to the base of the corresponding "tail"; the vesicles then probably play here also the rôle of receptacles for the "tails"; round refractile globules are also seen at the bases of the vesicles; the remainder of the body is filled by a homogeneous plasmic mass which frequently contracts to the center of the body cavity, forming a clearly distinct round or oval mass.

Habitat: Subcutaneous and superficial intermuscular tissue of *Coregonus fera* Jur.; April and May.

Myxobolus brevis Thélohan, 1892.

Henneguya brevis, Bull. Soc. philomat. Paris, IV, p. 177.

Myxobolus medius Thélohan, 1892.

Henneguya media Bull. Soc. philomat. Paris, IV, p. 176.

Myxobolus monurus, sp. nov.

(Psorosperms of *Aphredoderus sayanus* Ryder, 1880, Amer. Nat. XIV, pp. 211-212, figs. 1, 2.)

Cyst: Lenticular, large, about 20 in number.

Spore: Tailed; body lenticular or slightly obovate; tail undivided, thick at origin, gradually attenuating, more or less curved, 2 to 3 times as long as body; capsules 2, equal, subparallel.

Habitat: Subcutaneous intermuscular tissue of *A. sayanus* from near Woodbury, N. J.

Myxobolus macrurus, sp. nov.

Cyst: 15 to 20 or more, the size of a pinhead.

Spore: Plainly dorso-ventrally asymmetric; tailed; body round-elliptic; 10 or 11 μ long; 6 to 8 μ broad; 4 μ thick. Shell insoluble in sulphuric acid (cold, concentrated); superior valve more convex than inferior; tail somewhat less transparent than shell, completely dissolved by cold concentrated sulphuric acid; almost or entirely invisible in balsam; length 30 to 40 μ , usually much less, the attenuate posterior portion being easily, and consequently frequently, broken off; tail separated entirely from shell by iodine water. Capsules 2, equal; nuclei *ad plur.* 4, 2 of them being the pericornual; vacuole present.

Habitat: Cysts invariably† embedded in the subcutaneous tissue of some part of the head (especially the under surface of the lower jaw) of *Hybognathus nuchalis* Ag., from the Neches River, 14 miles east of Palestine, Texas, November 24, 1891.

* Dedicated to Dr. F. Zschokke, of Basel.

† Among several hundred cysts, one was seen at the base of the pectoral fin, say 3 mm. behind the head.

Myxobolus strongylurus, sp. nov.(Psorosperms of *Synodontis schal*, Müller, Müller's Archiv., 1841, pp. 480-481, t. 16, f. 2.)

Spore: Body anteriorly blunter than in *M. schizurus*; length of body 9 μ ; breadth 5.4 μ ; tail single, undivided, very peculiar in being constantly oblique in the longitudinal plane.

Habitat: Encysted in skin of head of *S. schal* from the Nile.

Myxobolus kolesnikovi,* sp. nov.(Psorosperms of *Coregonus fera* Kolesnikoff, 1886, Veter. Vestnik Kharkoff., v, pp. 242-248, f. 1-3.)

Cysts: Numerous (up to 80), spherical or oval, 10 to 30 mm. long by 7 to 20 mm. broad.

Spore: Round or oval with a sharp anterior end; tail single or double, thick at its origin, attenuating gradually.

Habitat: Interstitial connective tissue of the thoracic muscles of *Coregonus fera* Jur.

Kolesnikoff's figures show the "double" tail to be merely the separated (laterally shifted) halves of the really single tail. To this species should probably be approximated one of Claparède's 3 forms, † viz, the tailed form habitant in the muscles of *C. fera*.

Myxobolus linearis, sp. nov.(Psorosperms of *Pimelodus sebæ* and of *Platystoma fasciatum* Müller, Müller's Archiv., 1841, p. 489, t. 16, f. 10).

Spore: Body lanceolate-linear; length 3 to 4 times breadth; capsules 2, equal, entirely parallel-appressed; tail single, occasionally double.

Habitat: Cysts in membrane lining branchial cavity of *Rhamdia sebæ* Cuv. & Val., and in branchial lamellæ of *Pseudoplatystoma fasciatum* L., both from South American rivers.

In cysts at the base of the dorsal fin of *Ameiurus melas* Raf., from Storm Lake, Iowa, a spore occurs which I strongly suspect to be identical with this species, as it answers in every respect to the above (rather meager) diagnosis. It is peculiarly interesting, as the tail is composed of a dorsal and a ventral half, and is insoluble in sulphuric acid (cf. *M. macrurus*).

Myxobolus schizurus, sp. nov.(Psorosperms of *Esox lucius* Müller, Müller's Archiv., 1841, pp. 477-478, t. 16, f. 1.)

Cyst: 0.44 to 1.09 mm. in diameter.

Spore: Body oval, length 12 μ ; breadth 6 μ ; thickness one-half the breadth; tail stout at origin, 3 to 4 times length of body, very frequently (probably as a rule) more or less bifurcate; capsules 2, equal, diverging posteriorly.

Habitat: In cellular tissue of the eye muscles, in that of the sclerotic, and in that between the sclerotic and choroid of *Esox lucius* in May and June. Müller failed to find it in North American pikes.

* Dedicated to N. F. Kolesnikoff, who first figured this form.

† Claparède in Lunel's Hist. Nat. d. Poissons du bassin du Léman, pp. 113.

Myxobolus creplini,* sp. nov.

(Psorosperms of *Acerina vulgaris* Creplin., 1842, Wieg. Archiv. f. Naturgesch, 1, pp. 61-63, t. 1, f. 1 a-e; cf. Weltner, 1892, Sitzungs.-Ber. Ges. Naturf. Freunde Berlin, pp. 29-31, 34).

Spore: Body elongate ventricose-elliptic; length 17.3 μ ; breadth 5.8 μ ; tail simple, as long as or a little longer than the body; capsules 2, equal.

Habitat: On *Gymnocephalus cernua* L., collected March 14, 1837.

Weltner believes the form observed by him in the ovary of *Esox lucius* to be identical with this form. Weltner's species was sometimes tailed and sometimes untailed.

Myxobolus psorospermica Thélohan, 1892.

Henneguya psorospermica, Bull. Soc. philomat. Paris, IV, pp. 167, 176.

Myxobolus diplurus, sp. nov.

(Psorosperms of kidney of *Lota vulgaris* Bütschli, 1882, Bronn's Thier-Reich, 1, t. 38, f. 21.)

Description (from figure).

Spore: Capsules 2, equal, posteriorly situated; tail double from base, the 2 halves adnate.

Habitat: Kidney of *Lota lota* L.

CHLOROMYXIDÆ fam. nov.

("Chloromyxées" et "Myxidiées" (*pars*) Thélohan, 1892, Bull. Soc. philomat. Paris, IV, pp. 173, 176.)

Definition: *Phænocystes* destitute of antero-posterior, but possessing bilateral symmetry;† capsules in 1 group at the anterior end; a bivalve shell, the plane of junction of whose valves is perpendicular (?) to the longitudinal; no vacuole; type genus *Chloromyxum* Ming.

CHLOROMYXUM Mingazzini, 1890.

Bull. Soc. Nat. Napoli, IV, p. 160; *ib.*, *Sphærospora*, *Myxosoma* et *Mixosoma* † Thélohan, 1892, Bull. Soc. philomat. Paris, IV, pp. 173, 175-6.

Definition: *Chloromyxidæ* with subspherical or ovate spores, whose breadth does not exceed their length; valves hemispherical; sporoplasm bilaterally and symmetrically situated; type *C. leydigii*.

CHLOROMYXUM *sens. strict.*

Definition: Quadricapsulate *Chloromyxa*; type *C. leydigii*.

Chloromyxum fluviatile Thélohan, 1892.

Bull. Soc. philomat. Paris, IV, pp. 173, 176, fig. 2.

* Dedicated to the discoverer, J. C. L. Creplin.

† Imperfect from unilateral position of sporoplasm in *Ceratomyxa*.

‡ See subgenus *Sphærospora*, p. 419. The table on p. 411 shows that *Sphærospora* and *Mixosoma* differ from *Chloromyxum* by only a single character, viz: the number of the capsules. As shown on p. 411, this character is a subordinate one compared to the grouping and position of the capsules, in which latter all the three genera agree. They may, therefore, all be grouped under one genus.

The two unispecific genera *Sphærospora* and *Mixosoma* have (at least as far as the record now shows) absolutely no distinctive character but the shape of the spore. They are therefore fused. Provisionally (but with some hesitation) I have recognized *Sphærospora* (including *Mixosoma*) as a subgenus. Its sole claim to such distinction rests on 2 capsules as against 4 in *Chloromyxum* proper. It is also worthy of note that the possibility of transitions are by this definition arbitrarily excluded, inasmuch as all our experience shows that increase of capsule number is by duplication and not by addition. So that the possibility of its ultimate entire fusion with *Chloromyxum* seems by no means remote.

Chloromyxum mucronatum sp. nov.(Psorosperms of *Gadus lota* Lieberkühn, Müller's Archiv., 1854, pp. 352-3, 368, t. 14, f. 5, 6.)Myxosporidium: Measuring 75 μ or less; spherical or ellipsoidal, rarely irregular.Spore: Broadly rounded-oval; concave pointed anteriorly; length *ad max.*, 8 μ .Habitat: Free in urinary bladder of *Lota lota* L.**Chloromyxum leydigii** Mingaz., 1890.

Boll. Soc. Nat. Napoli., iv, pp. 160-4 (see also the next species).

Chloromyxum incisum sp. nov.(Psorosperms of *Raja batis* Leydig, Müller's Archiv., 1851, pp. 225-226, 234, t. 8, f. 4a-f.)Myxosporidium: Biliary yellow, roundish or somewhat elongate, 29 to 88 μ in diameter, without or with 1 to 4 pansporoblasts.Spore: Sharply cuneate-oval, *posterior border radiate-incised*, resembling a radiate-ribbed Lamellibranch shell.Habitat: Free in gall-bladder of *Raja batis* L.

In face of the striking difference between this spore-form and *C. leydigii*, the present evidence (which consists of Mingazzini's opinion* without any detailed evidence, Perugia's opinion,† too little explicit, and the probably not independent opinion of Thélohan ‡) is insufficient to warrant the fusion of the two forms, especially as it does not appear that either Mingazzini or Perugia examined the gall-bladder of *Raja batis*.

Chloromyxum ?? **congru** Perugia, 1891.*Myxosporidium congru*, Boll. Scientif. Pavia, XIII, pp. 24-5, f. 15-20.Habitat: Gall-bladder of *Leptocephalus conger* collected in August, 1890.

Subgenus SPHÆROSPORA Thélohan, 1892.

Bull. Soc. philomat. Paris, iv, p. 175; *Mixosoma*§ *et Myxosoma* *ibid.*, p. 175.Definition: Bicapsulate Chloromyxa; type *Chloromyxum* (*S.*) *elegans* Thél.**Chloromyxum elegans** Thélohan, 1892.*Sphærospora elegans*, Bull. Soc. philomat. Paris, iv, pp. 167, 175.**Chloromyxum dujardini** Thélohan, 1892.(Psorosperms of *Cyprinus rutilus* (*pars*) Müller, Müller's Archiv., 1841, pp. 481, 486, t. 16, f. 4 b, c; psorosperm of *Cyprinus erythrophthalmus* Dujardin, 1845, Hist. Nat. des Helminthes, p. 644, t. 12, f. 12 N₁, 12 N₂.)*Mixosoma dujardini*, Bull. Soc. philomat. Paris, iv, p. 175.

Myxosporidium: 1.25 to 1.50 mm. long.

Spore: Ovate, pointed anteriorly, broadly rounded posteriorly; length 10 to 12 μ ; capsules 2, of equal size.Habitat: Branchial lamellæ of *Leuciscus* (*Scardinius*) *erythrophthalmus* from the Vilaine, at Rennes, France; pseudobranchiæ of *Leuciscus rutilus*, from German rivers.

Concerning the form observed by him, Müller says:

Once there was found on the pseudobranchiæ (*Nebenkiemen*) a mass of small yellow cysts. The size of this mass was 4 lines. This time all the cysts contained elongate capsules with pointed anterior and bluntly rounded posterior ends (f. 4 b). On the flat border the convex surfaces were exactly equal and the two diverging vesicles were attached interiorly at their points.

* Boll. Soc. Nat. Napoli, 1890, iv, pp. 160-161.

† Boll. Scientif. Pavia, 1890, XII, p. 138.

‡ Bull. Soc. philomat. Paris, 1892, iv, p. 176.

§ Type *Chloromyxum* (*Mixosoma*) *dujardini*.

Thus this form was never found coexisting in the same cyst with *Myxobolus cycloides* sp. nov. Considering the great frequency of occurrence of the latter species such coexistence would be expected if they were merely different forms of one species. Their persistent non-association thus strongly reënforces the argument in favor of their specific distinctness, drawn from their different characters.

The synonymy is on the authority of Thélohan (letter to the author, 1893). He has found *Mixosoma dujardini* upon both of the fishes named above and believes that Müller's and Dujardin's figures represent the same species.

CERATOMYXA Thélohan, 1892.

Bull. Soc. philomat. Paris, iv, pp. 169, 175.

Definition (provisional): *Chloromyxidæ* with bilaterally symmetrical, transversely extended, sub-isosceles-triangular spores, whose breadth greatly exceeds the length; valves hollow-conical with solid tips; sporoplasm unilaterally and asymmetrically situated; type *C. sphærulosa* Thél.

Ceratomyxa sphærulosa Thélohan, 1892.

Bull. Soc. philomat. Paris, iv, pp. 171-3, 175, fig. 1.

MYXIDIIDÆ fam. nov.

(“Myxidiées” (*pars*) Thélohan, 1892, Bull. Soc. philomat. Paris, iv, pp. 173, 175.)

Definition (provisional, as regards negative characters): *Phænocystes* destitute of antero-posterior, but possessing bilateral symmetry; capsules in 2 groups in the (right and left) wings; no bivalve shell; no vacuole; type (and only) genus *Myxidium* Bütschli.

MYXIDIUM Bütschli, 1882.

Bronn's Thier-Reich, I, t. 38.

Definition: Characters those of the family; type *M. lieberkühnii* Bütschli.

Myxidium lieberkühnii Bütschli, 1882.

Bronn's Thier-Reich, I, t. 38, f. 12-15.

Probably a second species is Leydig's “psorosperm” of the bile-ducts of *Raja batis*, Müller's Archiv, 1851, pp. 226, 234, t. 8, f. 4g.

INDEX.

STREAMS AND LAKES OF WESTERN MONTANA AND NORTHWESTERN WYOMING.

	Page.		Page.
Annotated list of fishes obtained in Montana and Wyoming	41-52	Genera and species referred to—Continued.	
Aster Creek	20, 21	<i>Cervus canadensis</i>	55
Atlantic Creek	26, 29	<i>Coregonus williamsoni</i>	47
Beaver Dam Creek	30	<i>Cottus bairdi punctulatus</i>	28, 51, 52
Beaverhead River	31, 32	<i>Dibothrium cordiceps</i>	29
Big Blackfoot River	14	<i>Erethizon epixanthus</i>	56
Bitter Root River	18	<i>Eutænia sirtalis parietalis</i>	57
Blacktail Deer Creek	31	<i>Eutænia vagrans</i>	57
Bonner	14	<i>Gulo luscus</i>	56
Botteler's Springs	4, 39, 40	<i>Lagomys princeps</i>	55
Bozeman	40	<i>Leuciscus atrarius</i>	22, 23, 46
Bozeman Creek	33	<i>gilli</i>	18, 44, 45
Bridger Creek	33	<i>hydrophlox</i>	22, 44
Bridger Lake	29	<i>Limnæa palustris</i>	14, 18
Browns Gulch Creek	17, 18	<i>Lota lota maculosa</i>	31, 52
Child's Bed Rock drain	4, 35	<i>Margaritana margaritifera</i>	15
Clarke Fork of the Columbia	10, 11	<i>Mephitis</i>	56
Clarke Fork of the Columbia, streams tributary to	10-19	<i>Mylocheilus caurinus</i>	18
Classified list of waters examined	9, 10	<i>Ovis canadensis</i>	56
Climate of the region examined	8	<i>Ptychocheilus oregonensis</i>	18, 43
Columbia River basin	10-28	<i>Rana pipiens brachycephala</i>	57
Columbine Creek	30	<i>Rana pretiosa</i>	58
Cottonwood Creek	4, 14, 36	<i>Ranunculus</i>	31
Crawfish Creek	21	<i>Ranunculus aquatilis trichophyllus</i>	18, 37
Davies Springs	40	<i>Rhinichthys dulcis</i>	14, 17, 22, 36, 42, 43
Deer Lodge	36	<i>Salmo mykiss</i>	48-50
Deer Lodge River	15	<i>Salmo irideus</i>	52
Demersville	12	<i>Salmo trutta levenensis</i>	52
Dempsey Creek	16, 17	<i>Salmo fario</i>	52
Depew Creek	13, 37	<i>Salvelinus fontinalis</i>	52
Drainage of the region examined	9	<i>Salvelinus malma</i>	14, 50, 51
Firehole River	32, 54	<i>Sciurus richardsoni</i>	55
Fish-cultural station, requirements for	3, 4	<i>Tamias</i>	55
Flathead Lake	12	<i>Taxidea americana</i>	56
Flathead River	11, 12	<i>Thymallus signifer</i>	47, 48
Forests in and about Yellowstone Park, preservation of	59, 60	<i>Ursus americanus</i>	56
Gardiner River	54	Gibbon River	32, 54
Gardiner River, East Fork	55	Glenn Creek	38, 39
Genera and species referred to—		Hellgate River	39
<i>Amblystoma tigrinum</i>	11, 57	Horsethief Springs	4, 33, 37, 38
<i>Antilocapra americana</i>	56	Inness Lake	23
<i>Arctomys</i>	55	Itinerary, Brief statement of	4, 5
<i>Astacus gambelli</i>	21	Itinerary, Detailed statement of	6, 7
<i>Bison bison</i>	56	Jackson Lake	22
<i>Bufo halophilus</i>	57	Jay Creek	29
<i>Canis latrans</i>	56	Jocko River	11
<i>Cariacus macrotis</i>	56	Lewis fork of the Columbia, Streams tributary to	19, 23
<i>Castilleia</i>	17	Lewis Lake	20
<i>Castor canadensis</i>	55	Little Blackfoot River	14, 35, 36
<i>Catostomus ardens</i>	22, 42	Location of station, places examined with reference to	34-41
<i>catostomus</i>	11, 14, 17, 18, 36, 42	Lolo Creek	18, 37
<i>discobolus</i>	41	McCiellan Creek	34, 35
<i>macrochilus</i>	42	Madison River	32
		Mammals noticed in Montana and Wyoming	55, 56

STREAMS AND LAKES OF WESTERN MONTANA AND NORTHWESTERN WYOMING—Continued.

	Page.		Page.
Mission Creek.....	11	Persons referred to or quoted—Continued.	
Missoula.....	36, 37	Pratt, W. B.....	58
Missouri River Basin.....	29, 41	Reynolds, W. F.....	24
Mount Powell.....	17	Sargent, John D.....	22
Nez Perce Creek.....	54	Stejneger, Leonhard.....	57
Objects of the expedition.....	5	Test, Frederick C.....	57-59
Pacific Creek.....	23, 27	Traphagen, Frank.....	5, 16
Pelican Creek.....	30, 31	Walker, Robert C.....	5
Persons referred to or quoted:		Winstanley, E. A.....	5, 18
Albee, Mr.....	36	Physical features of the region examined.....	7-10
Anderson, George S.....	6	Places examined with reference to location of station.....	34-41
Beadle, W. B.....	58	Polecat Creek.....	21
Bickford, W. M.....	5, 37	President Camp.....	21, 22
Bielenberg, N. J.....	5, 16, 36	Prickly Pear Creek.....	34
Bridger, Jim.....	25	Pure water required for a station.....	3
Child, W. C.....	4, 5, 35	Race-track Creek.....	17
Chisholm, O. P.....	6	Rattlesnake Creek.....	4, 13, 36
Clapham, Burnside.....	4, 5, 12, 31, 39	Recommendations.....	60
Conley, Frank.....	5, 16	Red Rock River.....	31
Cooper, Walter.....	6	Reese Creek.....	39
Cope, E. D.....	58	Reptiles and batrachians collected, annotated list of.....	57-59
Cummins, R. R.....	6, 33, 37, 38	Requirements for successful fish-cultural station.....	3, 4
Davies, W. J.....	40	Rock Creek.....	16
Dutton, C. E.....	22, 31	Senecio Creek.....	29
Gannett, Henry.....	20	Shoshone Lake.....	19, 20
Gill, Theodore.....	45	Shoshone and Lewis lakes.....	54, 55
Hague, Arnold.....	8, 25, 29, 30, 59, 60	Snake River.....	19, 22
Harwood, Edward.....	4, 5, 6	Station, places examined with reference to locating.....	34-41
Hayden, F. V.....	7, 8, 24, 26, 33, 34, 59	Stocking waters of Yellowstone National Park with fish, results of.....	53-55
Hofer, Elwood.....	4, 21, 23, 25, 26, 32, 54, 55, 56	Summary of report.....	3, 6
Imes, W. A.....	6	Swan Lake.....	13
Jenkins, O. P.....	4, 5, 21, 39	Swan River.....	12, 13
Jones, W. A.....	24, 25	Thoroughfare Creek.....	29
Jordan, David Starr.....	30, 32	Tincup Joe.....	16
Lucas, E. R.....	31, 33, 55	Two-Ocean Pass.....	24-28
Merriam, C. Hart.....	20, 55, 58	Upper Yellowstone River.....	29
Morgan, Thomas.....	16	Waters examined, classified list of.....	9, 10
Muth, William.....	4, 5	Wolverton Spring.....	40, 41
Pitcher, Lieut.....	32, 54	Yount Peak.....	29
Power, T. C.....	5		

REPORT UPON INVESTIGATIONS MADE IN TEXAS IN 1891.

Acknowledgments.....	62, 63	Genera and species referred to—Continued,	
Big White Oak Bayou.....	69	Dorosoma cepedianum.....	68
Buffalo Bayou.....	69	Fundulus heteroclitus.....	84
Clear Creek.....	68	pallidus.....	84
Colorado River.....	73, 74	similis.....	84
Comal Springs.....	73	xenicus.....	85
Corpus Christi.....	71	Gambusia patruelis.....	68, 88
Crustacea collected, list of.....	89	Goniobasis alexandrensis.....	73
Dickinson Bayou.....	67, 68	Hippa ererita.....	89
Galveston Bay.....	64, 71	Hybognathus nuchalis.....	76
Galveston Packing Company.....	66	Hybopsis æstivalis marconis.....	82
Genera and species referred to—		Lepomis pallidus.....	68
Alpheus heterochaelis.....	90	Libinia dubia.....	89
Anodonta corpulenta.....	71	Limnæa desidiosa.....	20
Arenæus cribrarius.....	89	Lucania parva.....	87
Callinectes hastatus.....	89	Micropterus salmoides.....	70
Cambarus clarkii.....	90	Mollienesia latipinna.....	88
Campostoma anomalum.....	75	Notemigonus chrysoleucus.....	68, 84
Chænobryttus gulosus.....	68	Notropis cayuga atrocaudalis.....	76
Cliola vigilax.....	76	delectus.....	80
Cyprinodon variegatus.....	84	deliciosus.....	77
Dionda episcopa.....	75	fumeus.....	81

REPORT UPON INVESTIGATIONS MADE IN TEXAS IN 1891—Continued.

	Page.		Page.
Genera and species referred to—Continued.		Oysters in Galveston Bay	66
Notropis lutrensis	79	Persons referred to or quoted—	
nocomis	78	Bringhurst, George A	62
notemigonoides	81	Church, W. D	64
nux	77	Dana, F. L	62
swaini	79	Dumble, E. T	63
venustus	70, 79	Earl, R. E	87
Opsopœodus osculus	82	Frizell, Joseph P	74
Palæmonetes exilipes	90	Gilbert, Charles H	83
vulgaris	90	Grant, John A	63
Palæmon ohionis	90	Gurley, R. R	63, 76
Penæus brasiliensis	90	Hay, O. P	87
setiferus	90	Herndon, A. C	62
Panopeus depressus	89	Hettler, Joseph	62
herbstii	89	Jenkins, O. P	87
texanus	89	Jordan and Gilbert	80
Petrolisthes armatus	89	Kilper, Jacob	68
Phenacobius mirabilis	82	Landa, Joseph	73
Physa halei	72	Looscan, M	62
Planorbis bicarinatus	73	Lubbock, J. B	63
lentus	72	McDonald, John	63
liedmanni	72	McDonald, Marshall	61, 63
Sesarma cinerea	89	Nichols, Fred McC	62
Squilla empusa	90	Rathbun, Mary J	89
Tillandsia recurvata	72	Runge, Julius	62
usneoides	68, 69, 72	Russell, R. H	72
Tozeuma carolinensis	90	Scovell, J. T	63
Unio undulatus	72	Singley, J. A	63
Zygonectes escambie	87	Smith, H. M	85, 88
funduloides	85	Wilson, Harvey T. D	62
jenkinsi	86	Wilson, Robert E. C	62
notatus	87	Woolman, A. J	83
pulvereus	85	San Antonio River	71, 72
Givens Oyster Company	71	San Antonio Springs	72
Guadalupe River	72, 74	San Jacinto River	70
Houston	68	San Marcos River	73
Hunter Creek	69, 70	San Pedro Springs	72
Instructions as to character of station	61	Sims Bayou	69
Itinerary	63	South Galveston	67
Kownslar Place	68	Swan Lake	66
Long Lake	70, 71	Texan fishes, notes on	75-88
Neches River	71	Trinity River	70
Olmos Creek	72	Waters examined	63

REPORT ON THE FISHERIES OF THE GULF STATES.

Abbott, W. H	91	Shore industries of—	
Alabama, fisheries of	138-144	Alabama	144
Alligator industry	91, 112, 157	Florida	135-138
Alligator trade	137, 138	Louisiana	171
Canning industry	136, 154, 171, 184	Mississippi	153-154
Common and scientific names of products	99-102	Texas	184
Comparisons with 1880	107, 108	Shrimp canning	154, 171
Fishermen, nationalities of	91	Shrimp trade	171
Fish trade	135, 144, 154, 167, 184	Sponge trade	137
Florida, fisheries of	108-138	Stevenson, Charles H	91
Hall, Ansley	91	Tabular statements:	
Lobsters, planted in Galveston Bay	97	1. Value of fish and other products taken with each principal form of apparatus used in Gulf fisheries in 1890	96
Louisiana, fisheries of	155-171	2. Rigs of vessels employed in fisheries of Gulf region in 1890	97
Mississippi, fisheries of	145-154	3. Number of persons engaged in fisheries of Gulf States in 1890	104
Names of fishes, mollusks, etc.	99-102		
Oyster canning	136, 154, 169		
Oyster trade	135, 136, 144, 154, 171, 184		
Pound-net fishing, obstacles to	95		
Race, Edward E	95		

REPORT ON THE FISHERIES OF THE GULF STATES—Continued.

Tabular statements—Continued.	Page.	Tabular statements—Continued.	Page.
4. Nationality of persons engaged in fisheries of Gulf States in 1890	104	34. Showing by counties the apparatus and capital employed in Alabama fisheries in 1889 and 1890	141
5. Investment in fisheries of Gulf States in 1890	104	35. Showing by counties and species the yield of Alabama fisheries in 1889 and 1890	141
6. Products of fisheries of Gulf States in 1890	104-105	36. Showing by counties the yield of Alabama vessel fisheries in 1889 and 1890	142
7. Percentage of quantity and value of each species taken in fisheries of Gulf States in 1890	106	37. Showing by counties and species the yield of Alabama shore fisheries in 1889 and 1890	142
8. Comparative table showing extent of fisheries of Gulf States in 1880 and 1890	106	38. Showing by counties and apparatus the yield of Alabama shore fisheries in 1889 and 1890	143
9. Comparative table showing by States the oyster yield of Gulf States in 1880 and 1890	108	39. Wholesale oyster trade of Alabama in 1889 and 1890	144
10. Persons employed in fisheries of west coast of Florida	109	40. Wholesale fish trade of Alabama in 1889 and 1890	144
11. Nationality of persons engaged in fisheries of west coast of Florida	110	41. Persons employed in Mississippi fisheries	146
12. Apparatus and capital employed in fisheries of west coast of Florida	110	42. Nationality of persons employed in Mississippi fisheries	146
13. Products of fisheries of west coast of Florida	111	43. Apparatus and capital employed in Mississippi fisheries	146
14. Showing by counties the number of persons employed in fisheries of west coast of Florida in 1889 and 1890	112	44. Products of Mississippi fisheries	147
15. Showing by counties the apparatus and capital employed in fisheries of west coast of Florida in 1889 and 1890	113-115	45. Showing by counties the number of persons employed in Mississippi fisheries in 1889 and 1890	147
16. Showing by counties and species the yield of fisheries of west coast of Florida in 1889 and 1890	116-118	46. Showing by counties the number and value of vessels, boats, apparatus, etc., employed in Mississippi fisheries in 1889 and 1890	148
17. Showing by counties and species the yield of vessel fisheries of west coast of Florida in 1889 and 1890	119-120	47. Showing by counties the yield of Mississippi fisheries in 1889 and 1890	149
18. Showing by counties and species the yield of shore fisheries of west coast of Florida in 1889 and 1890	121-123	48. Showing by species and counties the yield of Mississippi vessel fisheries in 1889 and 1890	150
19. Summary by customs districts of vessel fisheries of west coast of Florida in 1889 and 1890	124	49. Showing by counties and species the yield of Mississippi shore fisheries in 1889 and 1890	151
20. Showing by customs districts and species the yield of vessel fisheries of west coast of Florida in 1889 and 1890	125	50. Showing by apparatus and species the yield of Mississippi shore fisheries in 1889 and 1890	152-153
21. Showing by counties and apparatus the yield of vessel fisheries of west coast of Florida in 1889 and 1890	126-128	51. Canning industry of Mississippi	154
22. Showing by counties and apparatus the yield of shore fisheries of west coast of Florida in 1889 and 1890	128-134	52. Showing by counties the wholesale oyster-packing trade of Mississippi in 1889 and 1890	154
23. Wholesale fish trade of west coast of Florida in 1889 and 1890	135	53. Wholesale fish trade of Mississippi in 1889 and 1890	154
24. Wholesale oyster trade of west coast of Florida in 1889 and 1890	136	54. Persons employed in Louisiana fisheries	156
25. Oyster-canning industry of west coast of Florida in 1889 and 1890	136	55. Showing the nationality of persons engaged in Louisiana fisheries in 1889 and 1890	156
26. Wholesale green-turtle trade of west coast of Florida in 1889 and 1890	136	56. Apparatus and capital employed in Louisiana fisheries	157
27. Wholesale sponge trade of west coast of Florida in 1889 and 1890	137	57. Products of Louisiana fisheries	157
28. Wholesale alligator trade of west coast of Florida in 1889 and 1890	138	58. Showing by parishes the number of persons employed in Louisiana fisheries in 1889 and 1890	158
29. Persons employed in Alabama fisheries	139	59. Showing by parishes the apparatus and capital employed in Louisiana fisheries in 1889 and 1890	158-160
30. Nationality of persons engaged in Alabama fisheries	139	60. Showing by parishes and species the yield of Louisiana fisheries in 1889 and 1890	160-162
31. Apparatus and capital employed in Alabama fisheries	139	61. Showing by parishes and species the yield of Louisiana vessel fisheries in 1889 and 1890	163
32. Products of Alabama fisheries	140	62. Showing by parishes and species the yield of Louisiana shore fisheries in 1889 and 1890	163-165
33. Showing by counties the number of persons employed in Alabama fisheries in 1889 and 1890	140	63. Showing by parishes and apparatus the yield of Louisiana vessel fisheries in 1889 and 1890	166-167
		64. Showing by parishes and apparatus the yield of Louisiana shore fisheries in 1889 and 1890	167-170
		65. Wholesale oyster trade and the canning of oysters and shrimp in Louisiana in 1889 and 1890	171
		66. Wholesale fresh-fish and shrimp trades of Louisiana in 1889 and 1890	171
		67. Persons employed in Texas fisheries	173

REPORT ON THE FISHERIES OF THE GULF STATES—Continued.

	Page.
Tabular statements—Continued.	
68. Nationality of persons engaged in the fisheries of Texas	173
69. Apparatus and capital employed in Texas fisheries	173
70. Products of Texas fisheries	174
71. Showing by counties the number of persons employed in Texas fisheries in 1889 and 1890....	174
72. Showing by counties the apparatus and capital employed in Texas fisheries in 1889 and 1890.....	175
73. Showing by counties and species the yield of Texas fisheries in 1889 and 1890.....	176-177

	Page.
Tabular statements—Continued.	
74. Showing by counties and species the yield of Texas vessel fisheries in 1889 and 1890.....	178
75. Showing by counties and species the yield of Texas shore fisheries in 1889 and 1890.....	179-180
76. Showing by counties and apparatus the yield of Texas shore fisheries in 1889 and 1890....	181-184
77. Wholesale oyster and fish trades and turtle canning of Texas in 1889 and 1890.....	184
Texas, fisheries of.....	170, 184
Turtle, canning	184
Turtle trade.....	136
Vessels, rigs of, represented in Gulf fisheries.....	96, 97

REPORT ON A COLLECTION OF FISHES FROM THE ALBEMARLE REGION OF NORTH CAROLINA.

Albemarle Sound.....	186
Common names of fishes of the Albemarle region, list of	188
Davis Bay.....	189
Economic fishes taken in 1890, tabular statement of...	186
Edenton.....	193
Edenton Bay	193-196
Elizabeth City	189
Genera and species referred to:	
<i>Achirus fasciatus</i>	196
<i>Acipenser sturio oxyrhynchus</i>	190, 193, 198
<i>Amia calva</i>	190, 193, 198
<i>Ameiurus albidus</i>	190, 193, 198
<i>nebulosus</i>	190, 194, 198
<i>Ælurichthys marinus</i>	194
<i>Anguilla chrysypa</i>	191, 195, 199
<i>Aphredoderus sayanus</i>	199
<i>Brevoortia tyrannus</i>	191, 195
<i>Cambarus blandingii</i>	193
<i>acuta</i>	197
Centrarchidæ	197
<i>Centrarchus macropterus</i>	199, 200
<i>Chænobryttus gulosus</i>	196, 199, 200
<i>Clupea æstivalis</i>	191, 195, 199
<i>mediocris</i>	195, 199
<i>pseudoharengus</i>	191, 195, 199
<i>sapidissima</i>	191, 195, 199
<i>Cyprinus carpio</i>	190, 199
<i>Dorosoma cepedianum</i>	191, 195, 199
<i>Enneacanthus obesus</i>	199
<i>simulans</i>	196
<i>Erimyzon sucetta</i>	194
<i>Etheostoma nigrum olmstedii</i>	192, 196, 200
<i>Fundulus diaphanus</i>	191, 195, 199
<i>Gambusia patruelis</i>	191, 193, 195, 199
<i>Hybognathus nuchalis</i>	190, 194, 199
<i>Lepomis auritus</i>	196
<i>gibbosus</i>	192, 196, 200
<i>pallidus</i>	196, 200
<i>Lucius americanus</i>	195, 199

Genera and species referred to—Continued.	
<i>Lucius reticulatus</i>	191, 195, 199
<i>Menidia beryllina</i>	192, 195
<i>Micropterus salmoides</i>	192, 196, 200
<i>Morone americana</i>	192, 196, 200
<i>Moxostoma anisurum</i>	198
<i>crassilabre</i>	194, 198
<i>Myriophyllum</i>	193, 196
<i>Notemigonus chrysoleucus</i>	191, 195, 199
<i>Notropis hudsonius</i>	190, 194
<i>niveus</i>	194
<i>whipplei</i>	194
<i>Palæmonetes exilipes</i>	189, 193
<i>Paralichthys lethostigma</i>	196, 200
<i>Perca flavescens</i>	192, 196, 200
<i>Pomoxis sparoides</i>	192, 196, 199
<i>Querimana gyrans</i>	192, 195, 199
<i>Roccus lineatus</i>	192, 196, 200
<i>Semotilus atromaculatus</i>	199
<i>Stizostedion vitreum</i>	192, 196, 200
<i>Tylosurus marinus</i>	191, 195
Introductory note	185-187
Pasquotank River	189-192
Persons referred to or quoted:	
Evermann, Barton W	194
Gilbert, Charles H.....	194
Jordan, David Starr	185, 187, 192, 194, 198
Kendall, W. C	192
Leary, J. L	194, 195
Rathbun, Richard	185
Skinner, H. G.....	195
Worth, S. G	198, 199, 200
Waters, George.....	190
Plymouth and vicinity	197
Reedy Point.....	193
Roanoke River	197-200
Specific names of fishes of the Albemarle region, list of.	187
Tabular statement of economic fishes taken in 1890....	186
Weldon.....	197

OBSERVATIONS ON THE SPAWNING HABITS OF THE SHAD.

Albemarle production	206
Chapman Point.....	204
Daily average temperature, influence of	206
Delaware River production.....	204, 205
Ferry Landing	204
Fort Washington seine production for 1887 and 1888....	205
Fort Washington seine, record of operations, 1887-1891.	202
Inequalities in production	202

Movements of shad controlled by temperature	205
Potency of creeks	205
Potomac River production	203
Shad-egg production of the Potomac River, 1888-1891..	204
Shad ova on Potomac River, cost of.....	201
Spawning region of the shad	201
Stony Point	203
Sutton Beach fishery	203

AQUATIC INVERTEBRATE FAUNA OF THE YELLOWSTONE NATIONAL PARK, WYOMING,
AND THE FLATHEAD REGION OF MONTANA.

	Page.		Page.
Aërial sound, note on.....	215	Genera and species referred to—Continued.	
Bridge Creek.....	227	Daphnia.....	214, 218, 227, 231
Bridge Bay.....	228	angulifera.....	218, 220
Discussion of the collections.....	213-214	arcuata.....	222
Descriptions of species and varieties.....	240-256	clathrata.....	235, 240-241
Duck Lake.....	230	dentata.....	244
Explanation of plates.....	258	dentifera.....	240, 243-244
Flathead Lake.....	236-238	pulex.....	214, 216, 218, 225, 226, 227, 228, 230, 232, 233, 237, 238, 242, 243
Flathead River system.....	236-239	pulex var. pulicaria.....	242-243
Genera and species referred to:		schœdleri.....	231, 232, 233, 234
Acroperus leucocephalus.....	222, 230, 232	thorata.....	237, 238, 239, 241, 242
Agabus.....	233, 234	Daphniidæ.....	235
Agrion.....	218, 220, 232, 233, 239	Deronectes.....	226, 228, 233
Agrioninæ.....	219, 230	griseostriatus.....	234
Alona.....	225, 230, 232, 239	Diptera.....	218
Allorchestes.....	216, 219, 221, 227, 235	Diaptomus.....	216, 217, 218, 219, 220, 221, 222, 225, 227, 228, 229, 233, 235, 238
dentata.....	214, 218, 219, 220, 225, 230, 232, 233, 238, 239	leptopus.....	253
inermis.....	225, 227, 228, 229	lintoni.....	216, 225, 230, 231, 232, 233, 234, 235, 252, 253
Amnicola.....	220, 229	piscinæ.....	232, 253, 254
Amphipoda.....	218	shoshone.....	214, 216, 219, 221, 225, 229, 232, 233, 235, 251-252
Annelida.....	233	sicilis.....	216, 219, 222, 225, 226, 229, 232
Aulostoma lacustris.....	218	stagnalis.....	251, 253
Bosmina.....	216, 227, 238	Diffugia.....	222
longirostris.....	222	globulosa.....	230
Cænis.....	222	Dytiscidæ.....	218, 232, 233, 238, 239
Candona.....	230	Echinopyxis.....	222
Ceriodaphnia.....	214, 227, 232, 233	Ephemera.....	220
reticulata.....	232, 233	Ephemeridæ.....	227, 239
Chætogaster.....	233	Epischura.....	220
Chara.....	226, 231	fluviatilis.....	254
Chironomus.....	214, 217, 218, 219, 220, 222, 225, 226, 227, 228, 231, 232, 233, 234, 235, 238, 239	lacustris.....	254, 255
Chydorus.....	216, 218, 231, 232	nevadensis columbiæ.....	238, 239, 254-255
sphæricus.....	225, 227, 229, 232, 233	nordenskiöldii.....	254
Cladocera.....	225, 233, 240, 244	Eurycercus.....	232, 238
Cladophora.....	215, 222	lamellatus.....	218, 222, 227, 239
Clepsine.....	220, 225, 226, 227, 231, 234, 235, 239	Gammarus.....	216, 217, 219, 221, 225, 226, 227, 230, 232, 233, 235, 238
elegans.....	220	robustus.....	216, 218, 219, 228, 229, 239
ornata.....	232	Graphoderes fasciaticollis.....	230
Cœlambus.....	219	Gregarinæ.....	216
Coleoptera.....	227, 239	Haliplus.....	228, 230, 235
Colymbetes.....	233	Holopedium gibberum.....	219, 222, 234
Conochilus.....	219, 220, 225, 227	Hydaticus.....	232
leptopus.....	225, 229, 256	Hydrachnidæ.....	218, 220, 228, 239
volvox.....	219	Hydra fusca.....	220, 222
Copepoda.....	233, 247, 255	Hydrophilidæ.....	238, 239
Corethra.....	214, 220, 228, 230, 233, 234	Hydrophilus.....	217, 220
Corisa.....	214, 217, 218, 219, 220, 226, 227, 228, 232, 233, 235, 239	Hydroporus.....	232, 233
Crustacea.....	238	Hygrotrechus.....	228, 231
Culex.....	218	Lacinularia socialis.....	214
Culicidæ.....	227	Leptodora.....	238
Cyclops.....	214, 216, 220, 225, 227, 231, 232, 233, 235, 238	hyalina.....	222, 239
capilliferus.....	248-249	Libellula.....	232
gyrinus.....	216, 218, 222, 225, 230, 238, 239	Libellulida.....	217
minnilus.....	216, 225, 230, 247	Libellulinæ.....	230
serratus.....	247, 248	Limnæa.....	227, 228, 229, 232, 233, 238, 239
serrulatus.....	216, 218, 225, 229	Macrothrix.....	227, 232
thomasi.....	238, 249-259	Melaniidæ.....	220
Cypris.....	216, 218, 225, 226, 227, 228, 232, 239	Monostyla.....	228
barbatus.....	227, 228, 244-246	cornuta.....	229
grandis.....	246		
Daphnella.....	220, 227		
brachyura.....	221, 222		

AQUATIC INVERTEBRATE FAUNA OF WYOMING AND MONTANA—Continued.

	Page.
Genera and species referred to—Continued.	
Monostyla ovata	256
Naidomorpha	230
Nelumbium	256
Nepheleis	226, 235, 239
4 striata	219, 220
maculata	218, 220, 221, 226, 228, 230, 232, 234, 239
obscura	216
Neuronia	217, 221
Notonecta	217, 218, 219, 231, 232, 233
Nuphar	218, 220
Oligochæta	219, 226
Oscillaria	229
Ostracoda	244-246
Paludinidæ	220
Physa	217, 218, 219, 220, 222, 226, 227, 228, 229, 233, 235, 238, 239
Phryganeidæ	216, 233
Pisidium	217, 218, 219, 220, 222, 226, 229, 231, 232, 234, 235, 238, 239
Planorbis	220, 229, 232, 233, 238, 239
exacutus	227
Plumatella	238, 239
Poduridæ	228, 230
Polyphemus	218, 227, 228
pediculus	216, 220, 222, 225, 228, 229, 232
Polyzoa	220
Potamogeton	215, 222
Pristina lacustris	238
Protozoa	256
Rotifera	256
Salmo mykiss	228
Scapholeberis	238
mucronatus	218, 225, 229, 230, 232
Sida	238
crystallina	220, 230, 231, 238, 239
Simocephalus	220
vetulus	227, 230, 231, 233
Simulium	228, 229
Sphærium	232
Spongilla	217, 226, 232, 234
fragilis	239

	Page.
Genera and species referred to—Continued.	
Stentor	230
igneus	230
var. fuliginosus	256
Stylaria lacustris	222
Tabanidæ	227
Turbellaria	233
Unionidæ	220
Valvatæ	220, 228
Vaucheria	222
Gardiner Lakelet	232
Gardiner River System	231-233
Grebe Lake	234-235
Heart Lake	220-222
Introductory	207
Lake of the Woods	230
Lewis Lake	218-220
Madison River System	234-235
Mary Lake	234
Persons referred to:	
Birge, E. A	244
Boutelle, F. A	210, 211
Brode, H. S	212
Evermann, B. W	207, 212, 213
Harwood, E. L	207
Hofer, Elwood	208, 211
Jordan, David S	207
Linton, Edwin	207, 208, 209, 218
Lucas, E. R	210, 231
McDonald, Marshall	207
Parchen, W. H	212
Pelican Creek	227
Shoshone Lake	214-218
Small ponds, collections from	232-233
Snake River System	214-222
Soda Butte Creek	229
Swan Lake	231-232, 239
Trip of 1890	208-211
Trip of 1891	212-213
Twin Lakes	231
Yellowstone River System	223-230
Yellowstone Lake	223-229

NOTES ON A COLLECTION OF FISHES FROM THE SOUTHERN TRIBUTARIES OF THE CUMBERLAND RIVER IN KENTUCKY AND TENNESSEE.

Beaver Creek	266
Big South Fork of the Cumberland River	266-268
Brimstone Creek	267
Canada Creek	267
Caney Fork River	262
Cumberland River	265
Eagle Creek	263
Little South Fork of the Cumberland River	266
New River	267
Obeys River	263-265
Otter Creek	266
Roaring River	263
Rock Creek	267
Round Lick Creek	261
Species enumerated in lists:	
Ambloplites rupestris	262, 263, 264, 268
Amblystoma punctatum	268
Ameiurus natalis	260, 261
nebulosus	264, 266
Aplodinotus grunniens	262, 265
Campostoma anomalum	260, 261, 262, 263, 264, 265, 266, 267

Species enumerated in lists—Continued.	
Chrosomus erythrogaster	263, 266
Catostomus nigricans	260, 262, 263, 264, 265, 266, 267
teres	326
Clupea chrysochloris	264, 265
Cottus bairdi	261, 263, 265, 268
Etheostoma aspro	262, 265, 268
blennioides	260, 261, 262, 263, 264, 266, 268
camurum	268
caprodes	260, 261, 262, 264, 265, 266, 268
cinereum	264, 268
cœruleum	263, 265, 266, 268
copelandi	265
evides	262, 264
macrocephalum	264, 268
obeyense	265, 266, 268
rufolineatum	260, 261, 262, 264, 268
simoterum atripinnis	260, 261, 262, 264, 268
squamiceps	260, 261
stigmæum	262, 265, 268
zonale	265

NOTES ON FISHES FROM THE SOUTHERN TRIBUTARIES OF THE CUMBERLAND RIVER—Continued.

	Page.
Species enumerated in lists—Continued.	
<i>Fundulus catenatus</i>	261, 264, 265, 266, 268
<i>Hybognathus nuchalis</i>	262
<i>Hybopsis amblops</i>	260, 261, 262, 264, 266, 267
<i>kentuckiensis</i>	260, 261, 262, 264, 266, 267
<i>storerianus</i>	262
<i>watauga</i>	262, 264
<i>Ictalurus punctatus</i>	262, 264
<i>Ictiobus difformis</i>	262, 265
<i>Labidesthes sicculus</i>	260, 261, 264
<i>Lagochila lacera</i>	260, 267
<i>Lepisosteus osseus</i>	262, 264, 265
<i>Leptops olivaris</i>	262
<i>Lepomis cyanellus</i>	266
<i>megalotis</i>	260, 261, 262, 263, 266, 268
<i>pallidus</i>	260, 261
<i>Micropterus dolomieu</i>	260, 262, 263, 264, 265, 266, 268
<i>salmoides</i>	261, 265, 268
<i>Macrolepidotum duquesnei</i>	260, 262, 264, 265, 266, 267
<i>Notropis ariommus</i>	261, 264, 267
<i>atherinoides</i>	261, 262, 264, 265, 267

	Page.
Species enumerated in lists—Continued.	
<i>Notropis boops</i>	258, 262
<i>galacturus</i>	260, 261, 262, 263, 264, 265, 266, 267
<i>heterodon</i>	267
<i>megalops</i>	260, 261, 262, 264, 265, 266, 267
<i>telescopus</i>	260, 264, 265, 267
<i>cycnocephalus</i>	261, 263, 264, 265, 266, 267
<i>whipplei</i>	260, 262, 264, 265, 266, 267
<i>Noturus flavus</i>	262, 265
<i>Phenacobius uranops</i>	264, 265
<i>Pimephales notatus</i>	260, 261, 262, 264, 266, 267
<i>Rhinichthys atronasus</i>	263, 265
<i>Semotilus atromaculatus</i>	261, 263, 266, 268
<i>Stizostedion vitreum</i>	262
<i>Zygonectes notatus</i>	260, 261
Spring Creek	260, 261, 263
Streams examined, list of	259
Stone River	259, 260
West Fork of the Stone River	259, 260
Willis Creek	265
Wolf River	263

REPORT ON THE FISHERIES OF THE SOUTH ATLANTIC STATES.

Alewife fishery	274, 355
Alligator industry	335, 343-345
Apparatus in South Atlantic fisheries	274-275, 277, 278, 295-304, 314-317, 326-330, 339-343, 350-353
Character of fisheries	274-275, 308-309, 334
Comparisons between 1880 and 1890	279-281
Fishery resources	272
Fishing-grounds	282-284, 307-308, 319, 333-334
Fish trade	305, 306, 331-332
Florida alligator industry	335, 343-345
description of coast and rivers	333-334
fisheries	333-345
by apparatus	339-343
counties	336-338
general statistics	335, 336
importance of fisheries	334
increase of fisheries	334
nature of fisheries	334
oyster canning	345
shore industries	345
Fresh-water fisheries	278, 320, 346-355
Geography of region	271, 282, 307, 319, 333-334
Georgia fisheries	319-332
by apparatus	326-330
counties	322-326
development of	319-321
importance of	319-321
fishing-grounds	319
general statistics	321-322
oyster canning	332
cultivation	320
investigations	320
surveys	320
trade	331, 332
shore industries	331-332
Importance of South Atlantic fisheries	274-275, 284, 308-309, 319-321, 334
List of fishes figured	356
Menhaden industry	305, 306
North Carolina fisheries	282-306
by apparatus	295-305

North Carolina fisheries, by counties	287-295
importance of	284
fishing-grounds	282
fish trade	305, 306
general statistics	285-286
geography of coast	228
importance of fisheries	284
menhaden industry	305, 306
oyster canning	305, 306
industry	305, 306
packing	305, 306
trade	305, 306
porpoise industry	305, 306
rank as fishing State	285
shore fishing industries	305-306
Oyster canning	305, 306, 318, 320, 332, 345
fishery	274, 287, 318, 320
investigations	320
packing	305, 306
planting	318, 320
surveys	284, 320
trade	305, 306, 318, 332, 345
Persons referred to or quoted—	
Abbott, W. H.	270
Drake, James C.	320
Earll, R. Edward	271, 334
Hall, Ansley	270
McDonald, Marshall	271
Race, E. E.	270
Ravenel, W. de C.	270, 318
Smith, Hugh M.	269
Stevenson, Charles H.	270
Winslow, Francis	284
Worth, S. G.	270
Porpoise fishery	274, 305
industry	305, 306
River basins, fisheries of	346-355
Salt-water fisheries	278, 320, 346
Shad fishery	274, 287, 295, 353-354
Shore fishing industries	305-306, 331-332, 345

REPORT ON THE FISHERIES OF THE SOUTH ATLANTIC STATES—Continued.

	Page.
South Carolina fisheries.....	307-318
by apparatus.....	314-317
counties.....	310-313
general statistics.....	309-310
oyster canning.....	318
planting.....	318
Sturgeon fishery.....	279, 355
Tabular statements:	
1. Persons employed in fisheries of South Atlantic States in 1890.....	276
2. Vessels, boats, apparatus, shore property, and cash capital employed in fisheries of South Atlantic States in 1890.....	276
3. Products of fisheries of South Atlantic States in 1890.....	277
4. Products, by apparatus, of fisheries of South Atlantic States in 1890.....	277
5. Values of different fisheries of South Atlantic States in 1890.....	278
6. Value of the salt-water and fresh-water fisheries of South Atlantic States in 1890.....	278
7. Comparative statement of number of persons engaged in fisheries of South Atlantic States in 1880 and 1890.....	280
8. Comparative statement of vessels, boats, apparatus, and property employed in fisheries of South Atlantic States in 1880 and 1890.....	280
9. Comparative statement of values of principal products of fisheries of South Atlantic States in 1880 and 1890.....	281
10. Persons employed in North Carolina fisheries in 1889 and 1890.....	286
11. Apparatus and capital employed in North Carolina fisheries in 1889 and 1890.....	286
12. Products of North Carolina fisheries in 1889 and 1890.....	286
13. Showing by counties the persons employed in North Carolina fisheries in 1889 and 1890.....	288
14. Showing by counties the apparatus and capital employed in North Carolina fisheries in 1889 and 1890.....	288-290
15. Showing by counties the yield of North Carolina shore fisheries in 1889 and 1890.....	291-294
16. Showing by counties and species the yield of North Carolina vessel fisheries in 1889 and 1890.....	295
17. Number of shad taken in each county in North Carolina in 1889 and 1890.....	295
18. Showing by counties, apparatus, and species the yield of North Carolina shore fisheries in 1889 and 1890.....	297-304
19. Porpoise industry of North Carolina in 1889 and 1890.....	306
20. Menhaden industry of North Carolina in 1889 and 1890.....	306
21. Wholesale fish trade of North Carolina in 1889 and 1890.....	306
22. Oyster industry of North Carolina in 1889 and 1890.....	306
23. Persons employed in South Carolina fisheries in 1889 and 1890.....	309
24. Apparatus and capital employed in South Carolina fisheries in 1889 and 1890.....	309
25. Products of South Carolina fisheries in 1889 and 1890.....	310
26. Showing by counties the persons employed in South Carolina fisheries in 1889 and 1890.....	310

	Page.
Tabular statements--Continued.	
27. Showing by counties the apparatus and capital employed in South Carolina fisheries in 1889 and 1890.....	311-312
28. Showing by counties and species the yield of South Carolina fisheries in 1889 and 1890.....	312-313
29. Showing by counties, apparatus, and species the yield of South Carolina shore fisheries in 1889 and 1890.....	314-317
30. Oyster canning industry of South Carolina in 1890.....	318
31. Persons employed in Georgia fisheries in 1889 and 1890.....	321
32. Apparatus and capital employed in Georgia fisheries in 1889 and 1890.....	322
33. Products of Georgia fisheries in 1889 and 1890.....	322
34. Showing by counties the number of persons employed in Georgia fisheries in 1889 and 1890.....	323
35. Showing by counties the apparatus and capital used in Georgia fisheries in 1889 and 1890.....	323-324
36. Showing by counties and species the yield of Georgia fisheries in 1889 and 1890.....	325-326
37. Showing by apparatus the yield of Georgia vessel fisheries in 1889 and 1890.....	326
38. Showing by counties, apparatus, and species the yield of Georgia shore fisheries in 1889 and 1890.....	327-330
39. Wholesale fish trade of Chatham County, Georgia, in 1890.....	331
40. Retail fish trade of Savannah, Ga., in 1890.....	332
41. Oyster-canning industry of Georgia in 1889 and 1890.....	332
42. Persons employed in fisheries of eastern Florida in 1889 and 1890.....	335
43. Apparatus and capital employed in fisheries of eastern Florida in 1889 and 1890.....	335
44. Products of fisheries of eastern Florida in 1889 and 1890.....	336
45. Showing by counties the number of persons employed in fisheries of eastern Florida in 1889 and 1890.....	336
46. Showing by counties the apparatus and capital employed in fisheries of eastern Florida in 1889 and 1890.....	337
47. Showing by counties and species the yield of fisheries of eastern Florida in 1889 and 1890.....	338
48. Showing by counties, apparatus, and species the yield of fisheries of eastern Florida in 1889 and 1890.....	339-343
49. Oyster-canning industry of Florida in 1889 and 1890.....	345
50. Showing by river basins the number of persons employed in fresh-water fisheries of South Atlantic States in 1889 and 1890.....	347
51. Showing by river basins the vessels, boats, apparatus, etc., employed in fresh-water fisheries of South Atlantic States in 1889 and 1890.....	347-348
52. Showing by river basins the products of fresh-water fisheries of South Atlantic States in 1889 and 1890.....	348-349
53. Showing by river basins and apparatus the products of fresh-water fisheries of South Atlantic States in 1889 and 1890.....	350-353
Vessel fisheries.....	274, 295
unimportance of.....	274, 287

REPORT ON THE EUROPEAN METHODS OF OYSTER-CULTURE.

	Page.		Page.
<i>Amphipleura</i> (<i>Navicula</i>) <i>fusiformis ostreana</i>	390	Leroux, Mr	390
Amrum	370, 373	Lijmfjord	379
Anderson, A.	405	Lisbon	367, 368
Arcachon	369, 371, 382, 384, 385, 394, 395	Lloregat Bay	366
Auray	371, 376, 379, 382, 385, 394, 395, 398	Lucrine Lake	358, 364, 365
Austin, G. L.	405	Marennes	376, 378, 385, 389, 390, 399
Belgium, oyster-culture in	389-392	Mare Piccolo	359, 360
Bergen-op-Zoom	384	Margate	393
Berrington, A. P.	405	McDonald, Marshall	357
Blackwater River	394, 400	Meinesz & Co	391
Blankenberghe	391	Milosa, Salvatore	363, 405
Bottemanne, C. J.	382, 387, 405	Möbius, Prof	370, 371, 373, 374, 375, 376, 377, 402, 405
Bouchon-Brandely, M.	363	Natural supply of oysters in England	393
Brénéguy	364	Newman, J.	405
Brightlingsea	394, 400	Newman, Mr	394, 400
Brindisi	359	Nieuport	391
Bruges	389	Ochtmann, Mr	384, 385, 386, 405
Burnham	394	Origin of oyster-culture	358
Cacilhas	367	Ossegor	375
Cancale	371, 399	Ostend	376, 386, 389, 390, 391, 392
Cape Finisterre	366	Oyster-culture in Belgium	389-392
Castello del Ovo	366	England	393-401
Colne River	394, 400	Germany	370-380
Conclusion	402-405	Holland	381-388
Coruña	366	Italy	358-366
Coste, M.	358, 364, 367	Spain and Portugal	366-370
Crouch River	394, 400	<i>Ostrea angulata</i>	367, 369, 402
Cultural processes in England	398-401	<i>edulis</i>	358, 402
Danish Cronicle	372	<i>edulis venetianæ</i>	358
Elbe River	380	<i>plicata</i>	358
Elevage in Holland	386, 387, 388	Oyster-land concessions in England	396, 397
England, oyster-culture in	393-401	Pollio, Domenico	365, 405
English oysters, grades of	400	Portugal and Spain, oyster-culture in	366-370
European system, summary of	406	Production in Holland	384, 385
Falmouth	400	Röm	370, 373
Faversham	399, 400	Sables d'Olonne	359
Feddersen, Messrs	405	San Martin de Noya Bay	366
Fishery orders in England	396	San Nicolas de Neda Bay	366
Föhr	373	Santa Luccia	366
Fryer, W.	405	Santander	366
Fusaro, Lake	358, 363, 364, 365	Schelde	381, 382, 383, 384, 386
Genoa, Gulf	358	Schleswig	379, 380
Germany, oyster-culture in	370-380	Sheppey, isle of	393
Goes	385, 386	Smith, J.	405
Grades of English oysters	400	Soudre	376
Graells, M.	367	Spain and Portugal, oyster-culture in	366-370
Hamman, M.	405	Stichert & Strache	389
Havesham	393	Storm, E.	405
Hayling Island	376, 395	Summary of European system	406
Herne Bay	399	Sylt	370, 373
Hoek, Dr.	373, 387, 395	Tagus River	367, 369
Holland, oyster-culture in	381-388	Tarente	358, 359, 363, 364, 371, 394
Holstein Bank	377	Tarentine Gulf	358, 363
Hubrecht, Prof.	404	Towse, T. Wrench	405
Husum	370, 371, 372, 373, 377, 378, 379	Trieste	358
Huxley, Prof.	394, 395, 401, 404	Van Zœlan, Baron Grœninx	386, 405
Introduction	357	Van Nisse, Mr	387
Italy, oyster-culture in	358, 366	Venice	358
Jerseke	369, 379	Ver Nieuwe, Dr. Anselme	391
Kentish Flats	393, 397	Wanklyn, A. C.	405
Kergurionné	374	Wattenmeer	370, 371, 373, 374, 375, 376, 377
Kiel	379, 380	Weinreich, Baurath	405
La Tremblade	376, 377, 378	Whitstable	369, 376, 393, 395, 396, 397, 398, 399
Lease of oyster-cultural property in Germany, abstract	380	Yvres River	389
Leasin'g oyster lands in Holland	387, 388	Zuyder Zee	381, 382

ON THE CLASSIFICATION OF THE MYXOSPORIDIA.

	Page.		Page.
<i>Acerina cernua</i> . (See <i>Gymnocephalus cernua</i> .)		<i>Myxobolus brevis</i>	416
<i>vulgaris</i> . (See <i>Gymnocephalus cernua</i> .)		<i>creplini</i>	418
<i>Alburnus alburnus</i> , <i>Myxobolus obesus</i> in	415	<i>cycloides</i>	415
<i>Ameiurus melas</i> , <i>Myxobolus sp. ad linearis affin.</i> in ..	417	<i>diplurus</i>	418
<i>Aphredoderus sayanus</i> , <i>Myxobolus monurus</i> in	416	<i>ellipsoides</i>	414
<i>Catostomus tuberculatus</i> . (See <i>Erimyzon sucetta</i> .)		<i>globosus</i>	415
<i>Ceratomyxa</i>	411, 412, 420	<i>inequalis</i>	414
<i>sphaerulosa</i>	420	<i>kolesnikovi</i>	417
<i>Chloromyxidæ</i>	412, 418	<i>linearis</i>	417
<i>Chloromyxum</i>	411, 412, 418	<i>lintoni</i>	414
?? <i>congru</i>	419	<i>macrurus</i>	416
<i>dujardini</i>	419	<i>medius</i>	416
<i>elegans</i>	419	<i>merlucii</i>	415
<i>fluviatile</i>	418	<i>monurus</i>	416
<i>incisum</i>	419	<i>mugilis</i>	414
<i>leydigii</i>	418, 419	<i>mülleri</i>	414
<i>mucronatum</i>	419	<i>obesus</i>	415
<i>Coregonus fera</i> , <i>Myxobolus kolesnikovi</i> in	417	<i>oblongus</i>	414
<i>Myxobolus spheralis</i> in	415	<i>oviformis</i>	414
<i>Myxobolus ?? zschokkei</i> in	416	<i>perlatus</i>	415
<i>Cottus scorpio</i> , <i>Pleistophora typicalis</i> in	410	<i>piriformis</i>	414
<i>Cryptocystes</i>	409	<i>psorospermica</i>	418
<i>Cystodiscidæ</i>	412, 413	<i>schizurus</i>	417
<i>Cystodiscus</i>	411, 412, 413	<i>spheralis</i>	415
? <i>diploxys</i>	411, 413	<i>strongylurus</i>	417
<i>immersus</i>	413	<i>transovalis</i>	415
<i>Cyprinodon variegatus</i> , <i>Myxobolus lintoni</i> in	414	<i>unicapsulatus</i>	414
<i>Cyprinus erythrophthalmus</i> . (See <i>Leuciscus (S.) ery-</i>		?? <i>zschokkei</i>	416
<i>throphthalmus</i> .)		<i>Myxosoma</i> . (See <i>Mixosoma</i> .)	
<i>rutilus</i> . (See <i>Leuciscus rutilus</i> .)		<i>Myxosporidiæ</i> , synonym for <i>Myxobolidæ</i>	413
<i>Erimyzon sucetta</i> , <i>Myxobolus oblongus</i> in	414	<i>Myxosporidium</i> , synonym for <i>Myxobolus</i>	413
<i>Myxobolus globosus</i> in	415	<i>merlucii</i> (see <i>Myxobolus merlucii</i>) ..	415
<i>Esox lucius</i> , <i>Myxobolus schizurus</i> in	417	<i>mugilis</i> (see <i>Myxobolus mugilis</i>)	414
<i>Gadus lota</i> . (See <i>Lota lota</i> .)		<i>Palæmonetes varians</i> , <i>Thélohania macrocystis</i> in	410
<i>Glugea</i>	409	<i>Phænocystes</i>	409, 410
<i>anomala</i>	409	<i>Phoxinus funduloides</i> , <i>Myxobolus transovalis</i> in	415
<i>destruens</i>	409	<i>Pimelodus blochii</i> . (See <i>P. clarias</i> .)	
<i>microspora</i> (synonym for <i>anomala</i>)	409	<i>clarias</i> , <i>Myxobolus inequalis</i> in	414
<i>Glugeidæ</i>	409	<i>sebæ</i> . (See <i>Rhamdia sebæ</i> .)	
<i>Gymnocephalus cernua</i> , <i>Myxobolus creplini</i> in	418	<i>Platystoma fasciatum</i> . (See <i>Pseudoplatystoma fascia-</i>	
<i>Myxobolus perlatus</i> in	418	<i>tum</i> .)	
<i>Henneguya</i> (synonym for <i>Myxobolus</i>)	411, 412, 413	<i>Pleistophora</i>	409, 410
<i>Hybognathus nuchalis</i> , <i>Myxobolus macrurus</i> in	416	<i>typicalis</i>	410
<i>Labeo niloticus</i> , <i>Myxobolus unicapsulatus</i> in	414	<i>Pseudoplatystoma fasciatum</i> , <i>Myxobolus linearis</i> in ..	417
<i>Leptocephalus conger</i> , <i>Chloromyxum ?? congru</i> in	419	<i>Pyralis viridana</i> . (See <i>Tortrix viridana</i> .)	
<i>Lota lota</i> , <i>Myxobolus diplurus</i> in	418	<i>Raja batis</i> , <i>Chloromyxum incisum</i> in	419
<i>Chloromyxum mucronatum</i> in	419	<i>Rhamdia sebæ</i> , <i>Myxobolus linearis</i> in	417
<i>Lota vulgaris</i> . (See <i>L. lota</i> .)		<i>Silurus clarias</i> . (See <i>Pimelodus clarias</i> .)	
<i>Leuciscus rutilus</i> , <i>Chloromyxum dujardini</i> in	419	<i>Sphaerospora</i> , subgenus of <i>Chloromyxum</i> ...	411, 412, 418, 419
<i>Myxobolus cycloides</i> in	415	<i>Synodon hal</i> , <i>Myxobolus strongylurus</i> in	417
<i>erythrophthalmus</i> , <i>Chloromyxum dujardini</i> in ..	419	<i>Thélob</i>	409, 410
<i>Mixosoma</i> (synonym for <i>Chloromyxum</i>)	411, 412, 418, 419	<i>contejeani</i>	410
<i>Myxidiidæ</i>	412, 420	<i>giardi</i>	410
<i>Myxidium</i>	411, 412, 420	<i>macrocystis</i>	410
<i>lieberkühni</i>	410, 420	<i>octospora</i>	410
? <i>sp</i>	420	<i>Tinca tinca</i> , <i>Myxobolus bicostatus</i> in	414
<i>Myxobolidæ</i>	412	<i>vulgaris</i> . (See <i>T. tinca</i> .)	
<i>Myxobolus</i>	411, 412, 413	<i>Tortrix viridana</i> , <i>Cystodiscus ? diploxys</i> in	413
<i>bicostatus</i>	414		